

## Considerations about Traumatic Diaphragmatic Rupture, Case Series and Review of Literature

**Agron Dogjani<sup>1\*</sup>, Henri Kolani<sup>2</sup>, Kastriot Haxhirexha<sup>3</sup>, Arben Gjata<sup>4</sup>, Edvin Selmani<sup>1,4</sup>, Bleriana Shqepa<sup>5</sup> and Hysni Bendo<sup>5</sup>**

<sup>1</sup>University Hospital of Trauma, Lecturer at University of Medicine, Tirana, ALBANIA.

<sup>2</sup>Department of General Surgery, University Hospital Center "Mother Theresa", Tirana, ALBANIA.

<sup>3</sup>General Surgeon at Clinical Hospital of Tetovo, Tetovo, RN of MACEDONIA

<sup>4</sup>University of Medicine of Tirana, ALBANIA.

<sup>5</sup>Resident at University of Medicine of Tirana, ALBANIA.

**\*Corresponding author:** Agron Dogjani, University Hospital of Trauma, Lecturer at University of Medicine, Tirana, Albania

**Citation:** Dogjani A, Kolani H, Haxhirexha K, Gjata A, Selmani E, et al. (2020) Considerations about Traumatic Diaphragmatic Rupture, Case Series and Review of Literature. J Surg 5: 1315. DOI: 10.29011/2575-9760.001315

**Received Date:** 16 June, 2020; **Accepted Date:** 26 June, 2020; **Published Date:** 30 June, 2020

### Abstract

**Introduction:** Diaphragmatic rupture following trauma is often associated and missed injury. Traumatic diaphragmatic rupture remains a diagnostic challenge, and associated injuries determine the outcome in those cases diagnosed early. Penetrating trauma with direct injury to the diaphragm is more common and accounts for about two-thirds of cases. The incidence of injury is 5% in the thoracic-abdominal trauma [1-3]. This report documents our experience of treating traumatic diaphragmatic rupture in our Level 1 University Trauma Center.

**Methods:** We retrospectively analyzed 18 patients diagnosed with traumatic diaphragmatic rupture was presented in the Emergency Department from January 2019 to June 2019 caused by blunt trauma in 8 and by penetrating trauma in 4 cases.

**Results:** In our study were included 11 patients, 10 (90.9%) males, and 1(9.1%) female. The average age is 48.5 years, we have the age group 17-60 years where we have 9 (81.8%) cases, The diagnosis was made by chest X-ray in 7(72.7%), thorax and upper abdominal computed tomography in 45.40% of cases. The most common injured organs were the spleen (54.50%), liver (36.40%), Big Blood Vessels (27.3%), and intestinal (9.10%). The mortality rate was 18%.

**Conclusions:** Early diagnosis and early surgical treatment determine successful management of TDR, with or without herniation of the abdominal organs. Penetrating trauma with direct injury to the diaphragm is more common. Traumatic injuries of the left side are more frequent, consequently, the most damaged organ is the spleen.

**Keywords:** Diaphragm; Intervention; Postsurgical complications; Rupture Of Stay; TDI: Traumatic Diaphragmatic Injury; ED: Emergency Department; OM: Operator Management

**Abbreviations:** TDR: Traumatic Diaphragmatic Rupture; MVA: Motor Vehicle Accident; FFH: Fall From Height; HWSO: Hit With Strong Object; GSW: Gunshot Wound; SW: Stab Wound; STW: Sharp Tools Wound; EP: Tromboembolism Pulmonary; MOF: Multi Organ Failure; DIC: Disseminated Intravascular Coagulation; MRI: Magnetic Resonance Imaging; CT: Computerized Tomography Scanner; DPL: Diagnostic Peritoneal Lavage; BI: Blast Injuries; BT: Blunt Trauma; PT: Penetrant Trauma; SBP: Systolic Blood Pressure; HR: Heart Rate; HDCO: Hemodynamic Condition; FAST: Focused Assessment With Sonography In Trauma; ICU: Intensive Care Unit; LOS: Length

### Introduction

The Diaphragm is the main respiratory muscle and the second most important muscle after the heart, it is so important for the respiratory process that any disorder of its function or damage to it is vital to the body [2]. Diaphragmatic injuries are relatively rare and may be due to Blunt (abdominal or thoracic) trauma, penetrating trauma, or a combination of trauma [1]. TDR (Traumatic Diaphragmatic Rupture), occurs in 5% of patients with thoracic-abdominal trauma and abdominal mass herniation occurs in 45-60% of patients. In recent decades, the treatment of TDR

has moved significantly [3]. The incidence of TDR has increased in recent years, parallel to the increase in the frequency of abdominal injuries, caused by MVA (Motor Vehicle Accidents) or occupational accidents [1-4]. In establishing a more accurate and rapid diagnosis in addition to evaluate the mechanism of injury, TDR requires careful evaluation of other diagnostic methods such as Chest x-ray, CT, MRI (Magnetic Resonance imaging), [3] or even the use of laparoscopy and DPL [5]. The objectives of this article are: to identify epidemiological data, the mechanism of damage for both types of TDR, to evaluate pre-hospital management as an important element in the early diagnosis of diaphragmatic trauma or concomitant injuries, or the localization of the injury and the correlation between it and the time of intervention by comparing it with the data of the literature, to illustrate postoperative complications according to Clavien - Dindo classification and their distribution and correlation with ISS.

## Material and Method

Our study was retrospective, conducted within the period of January 2019 to June 2019. Our study included 11 patients who were presented at our University Trauma Hospital. In our study, we were based on a few points that will make it easy: Grouping of patients by gender and age, from sort of trauma according to the trauma mechanism; in accordance with the time of hospitalization; under the transport mode; by ISS value; by the time of Surgery; with the accompanying damaged organs; to the postoperative complications; from the cause of death. Discrete data analysis (variables) were presented in absolute value and percentage.

## Results

In our study were included 11 patients, 10(90.9%) males, and 1(9.1%) female. The average age is 48.5 years, we had most affected the age group 17-60 years where we have 9(81.8%) of cases. According to the etiology of TDR; 8(72.7%) cases were after Penetrant Trauma (PT), and 3(27.3%) cases from Blunt Trauma (BT). Regarding the mechanism of injury, most of them were after penetrating injuries with 8(72.7%) cases, of which the most common were stab wounds (SW) in 4(50.0%) cases, followed by Gunshot Wounds (GSW) with 2(25.0%), while injuries with blast injuries(BI) and with Sharp Tools Wounds (STW) with one case. While after blunt injuries were 3(27.3%) cases, and 2(33.3%) cases after the Motor Vehicle Accident (MVA) and one case after the Fall from Height (FfH). All trauma patients arrived at the hospital during the first 6 hours.

The patient's condition (hemodynamically condition - HdCo) in arrival based on the Allgower Formula (rate of SBP/HR) in most of cases were potentially risked in 6(55%) of cases, followed with stable hemodynamically in 4(36%) of cases and with hemorrhagic shock in one case. Most of trauma patients (64%) were transported with ambulance, and 36% of cases with private cars. The value

of ISS at our patients had an average of 23.7. The performance of the examinations as part of the primary and secondary survey to do early diagnosis in diaphragm injury was Chest x-ray in 7(72.7%), Abdomen x-ray in 2(18.20%), CT in 5(45.40%), FAST in 6(54.50%) of cases. According to the side of diaphragm injuries, we have found the left side in 8(73%), and to right side in 3(27%) of cases. All cases had OM and the time of Surgery was in most of cases within the first 2 hours to 8(72, 7%), after 2-6 hours to 2(18.20%), and after 6 hours with one case (Table 1).

| Surgery Time | Localisation / of Rupture |                 | Total           |
|--------------|---------------------------|-----------------|-----------------|
|              | Left Side                 | Right Side      |                 |
| 0-2h         | 7(63.6%)                  | 1(9.1%)         | 8(72.7%)        |
| After 2-6h   | 1(9.1%)                   | 1(9.1%)         | 2(18.2%)        |
| After 6-24h  | 0(0%)                     | 1(9.1%)         | 1(9.1%)         |
| <b>Total</b> | <b>8(72.7%)</b>           | <b>3(27.3%)</b> | <b>11(100%)</b> |

**Table 1:** Distribution of DTR in according the Surgery Time.

The accompanying injuries were as follows; Spleen in 6(54.5%), ribs in 5(45.50%), chest 4(45.5%), liver 4(36.5%), big blood vessels injury 2(18.20%), intestinal and pericardial injury in one case (Table 2).

| Accompanying Injury | References from Literature | Our Results |
|---------------------|----------------------------|-------------|
| Spleen              | 50%                        | 54.50%      |
| Ribs                | 47.20%                     | 45.50%      |
| Liver               | 38.60%                     | 36.40%      |
| Lungs               | 30.90%                     | 27.3%       |
| Big Blood Vessels   | 11.1%                      | 18.20%      |
| Intestinal          | 19.40%                     | 9.10%       |
| Pericardia          | 8.0%                       | 9.10%       |

**Table 2:** Distribution of accompanying injuries in DTR.

During TDR treatment, it was also associated with other surgical procedures, with this distribution; Chest drainage at 9(90.9%) of cases; Liver suture in 3(36.4%) of cases; Splenectomy in 3(27.3%) of cases; Lien preservation in 1(9.1%) case, Primary Vascular Suture, Pericardial Suture and primary Stomach suture in one case. The recovery period was in most of the cases to 6(64%) in the surgical ward, and in 3(36%) of cases were treated in ICU. Evaluation of surgical wounds was per primam (not infected) in 9(82%) of cases and infected wounds in 2(18%) of cases. Length Of Stay (LOS) in hospital was; more than 7 days in 10(90.0 %), and 0-1 day in one case; The distribution of Post-operative complications was: Pleural Versament 36.4%, Sepsis 27.3%,

Hemopneumothorax, Eventration, Biliary Fistula, Respiratory Acute insufficiency with one case. Mortality in our cases was 18%, while the time when it happened was one case during the surgery and one case after 72 hours. The causes of death, was result of fibrinolysis in one case, and Thromboembolic of Pulmonary (TEP) in one case.

## Discussion

In our study, we included 11 patients, 9.1% female, and 90.9% male, whose average age was 48.5 years. These data are in line with those in the literature Hofmann, Sabine, et al [6], and the values being as follows: males 88.2%, females 11.8% and median age 46 years. We also have a match with age distribution. In our study we showed that 82% were in the 17- 60 years age group and 18% in the 61- 60 years age group, whereas in the literature Gwely, et al [5], this distribution is described as 83 % for 17-60 years and 10.9% over 61 years. The most common etiology of TDR has penetrated injury to 8(73%) of cases, followed by blunt injury with 3 (27%) of cases. Even in this case, we have a concordance with the literature data where TP causes 73% of cases with TDR [7,8]. Zarour, et al. [8] report the largest single-institution series in the current literature of 773 patients with TDI over a 15-year period. The majority (73%) of these patients were penetrating trauma victims, and most were managed with laparotomy and direct suture repair of the injury.

Every patient with penetrating trauma should be examined for diaphragmatic damage. In terms of mechanism of injury, most of them were after penetrating injuries with 8(72.7%) cases, of which the most common were stab wounds (SW) in 4(50.0%) cases, followed by Gunshot Wounds (GSW) with 2(25.0%), while injuries with blast injuries(BI) and with Sharp Tools Wounds (STW) with one case. While after blunt injuries were 3 (27.3%) cases, and 2(33.3%) cases after the Motor Vehicle Accident (MVA) and one case after the fall from Height (FfH). All trauma patients arrived at the hospital during the first 6 hours. The time of hospitalization, which is an important point, turns out to be 100% in the first 6 hours. The shorter time to submit to ED means a shorter time to diagnose TDR, faster OM, fewer complications from the herniation, and ultimately a reduction in mortality and morbidity. The proximity of our Center with localization in 81.8% of cases and the proximity of national roads to hospitals are the reasons why we have this distribution. These results are in agreement with the literature, Beng Leong Lim, et al [2]. Tseng J, et al [9]. "The Allgower Formula or Shock Index is reported to carry significant prognostic value for trauma and other emergency cases admitted to the emergency department".

In our study the patient's condition (hemodynamically condition - HdCo) in arrival based on the Allgower Formula

(rate of SBP/HR) in most of the cases were potentially risked in 6(55%) of cases, followed with stable hemodynamically in 4 (36%) of cases and with hemorrhagic shock in one case. The ISS has a gravity injury rate of 23.7 for those diagnosed with TDR and values below 16 represent 3(27.3%) of cases, values from 16-26 represent 5(45.5%) of cases and those above 27 represent 3(27.3%) of cases. These results are consistent with recent years' studies from Gwely, et al [10]. According to the site of injuries, we have found: the left side in 7(73%) of cases and the right side in 27% of cases. Right diaphragmatic injuries, especially after BT, are relatively rare and have a relative loss/delay of diagnosis and consequently delayed surgical treatment [11]. Missed diagnosis of a large, right-sided diaphragmatic rupture with herniated liver and concomitant liver laceration after blunt trauma: consequences for delayed surgical repair. However, many authors believe that TDR right side is associated with higher mortality, so they remain undiagnosed while undergoing autopsy. [10,12,13]. Mortality in our cases was 18%, while the time when it happened was one case during the surgery and one case after 72 hours. The causes of death were result of fibrinolysis in one case, and Thromboembolic of Pulmonary (TEP) in one case. The accompanying injuries were as follows: Spleen in 6(54.5%), ribs in 5(45.5%), chest 4 (45.5%), liver 4(36.5%), big blood vessels injury 2(18.20%), intestinal and pericardial injury in one case (Table 2). So, given the limited number of cases we had in the study, we find that we have a concordance with the literature data on the most frequent organ injuries: spleen, Liver, Lung [1,8]. The adult ISS, right from the patient's first presentation, is rightly proportional to either morbidity or mortality, and even postoperative complications are proportional to them [3]. Mortality in our study was 18% and survival was 82%. In the literature, mortality is described by 10% -35%, so we have a good agreement with our study [4,14-18].

## Conclusion

Early diagnosis and early surgical treatment determine successful management of TDR, with or without herniation of the abdominal organs. Penetrating trauma with direct injury to the diaphragm is more common. Traumatic injuries of the left side are more frequent, consequently, the most damaged organ is the spleen.

**Acknowledgements:** the authors would like to thank Department of General Surgery and Head of the Department Surgery for providing support in preparing the manuscript.

**Contributions:** all authors contributed to the study conception and design of the study and approved the final version of the manuscript. AD analyzed and interpreted data; KH, ES, AGj drafted the article and reference search; HK, HB, BSh did the manuscript reviewing.

**COI Statement:** This paper has not been submitted in parallel. It has not been presented fully or partially at a meeting or podium or congress. It has not been published nor submitted for consideration beforehand.

All authors declare that there is no conflict of interest. This research received no specific grant from any funding agency in the public, commercial, or nonprofit sectors. There are no relevant or minor financial relationships from authors, their relatives or next of kin with external companies.

**Disclosure:** The authors declared no conflict of interest. No funding was received for this study.

## References

1. Al-Refaie RE, Awad E, Mokbel EM (2009) Blunt traumatic diaphragmatic rupture: a retrospective observational study of 46 patients. *InteractCardiovascThoracSurg* 9: 45-49.
2. Beng Leong Lim, Li Tserng Teo, Ming Terk Chiu, Marxengel L Asinas-Tan, Eillyne Seow (2017) Traumatic diaphragmatic injuries: a retrospective review of a 12-year experience at a tertiary trauma centre. *Singapore Med J* 58: 595-600.
3. Bodanapally UK, Shanmuganathan K, Mirvis SE, Clint W Sliker, Thorsten R Fleiter, et al. (2009) MDCT diagnosis of penetrating diaphragm injury. *Eur Radiol* 19: 1875-1881.
4. Chugtai T, Syed Ali, Sharkey P, Lins M, Rizoli S (2009) Update on managing diaphragmatic rupture in blunt trauma: a review of 208 consecutive cases. *Can JSurg* 52: 177-181.
5. Noureldin Noaman G (2010) □ Outcome of blunt diaphragmatic rupture. Analysis of 44 cases. □ *Asian cardiovascular & thoracic annals* 18: 240-243.
6. Sabine Hofmann, Marko Kornmann, Doris Henne-Bruns, Andrea Formentini (2012) □ Traumatic diaphragmatic ruptures: clinical presentation, diagnosis and surgical approach in adults. □ *GMS Interdisciplinary plastic and reconstructive surgery DGPW* 2012.
7. Lopez PP, Arango J, Gallup TM, Stephen M Cohn, John Myers et al. (2010) Diaphragmatic injuries: what has changed over a 20-year period? *Am Surg* 76: 512-516.
8. Zarour AM, El-Menyar A, Al-Thani H, Scalea TM, Chiu WC (2013) Presentations and outcomes in patients with traumatic diaphragmatic injury: a 15-year experience. *J Trauma Acute Care Surg* 74:1392-1398.
9. Tseng J, Nugent K (2015) Utility of the shock index in patients with trauma. *Am J Med Sci* 349: 531-535.
10. Hanna WC, Ferri LE, Fata P, Razek T, Mulder DS (2008) The current status of traumatic diaphragmatic injury: lessons learned from 105 patients over 13 years. *Ann Thorac Surg* 85: 1044-1048.
11. Simon LV, Burns B (2019) Diaphragm Rupture. *State Pearls* [Internet] 2019.
12. Hanna WC, Ferri LE (2009) Acute traumatic diaphragmatic injury. *Thorac Surg Clin* 19: 485-489.
13. Rashid F, Chakrabarty M, Singh R, Iftikhar S (2009) A review on delayed presentation of diaphragmatic rupture. *World J Emerg Surg* 4: 32.
14. Clavien PA, Barkun J, de Oliveira ML, Jean Nicolas Vauthey, Daniel Dindo, et al. (2009) The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg* 250: 187-196.
15. Demetriades D, Kakoyiannis S, Parekh D, Hatzitheofilou C (1988) Penetrating injuries of the diaphragm. *Br J Surg* 75: 824-826.
16. Morgan BS, Watcyn-Jones T, Garner JP (2010) Traumatic Diaphragmatic Injury. *JR Army Med Corps* 156: 139-144.
17. Murray JG, Caoili E, Gruden JF, Evans SJ, Halvorsen RA, et al. (1996) Acute rupture of the diaphragm due to blunt trauma: diagnostic sensitivity and specificity of CT. *AJR Am J Roentgenol* 166: 1035-1039.
18. Søreide K, Reite A, Haaverstad R (2017) Missed diagnosis of a large, right-sided diaphragmatic rupture with herniated liver and concomitant liver laceration after blunt trauma: consequences for delayed surgical repair. *J Surg Case Rep* 2017: 157.