

Research Article

Emergency Department Approach to the Growing Global Public Health Problem: Violence Against Women

Ali Kemal Erenler^{1*}, Mehmet Oğuzhan Ay¹, Ahmet Baydın²

¹Department of Emergency Medicine Hıtit University Çorum Education and Research Hospital, Çorum, Turkey

²Department of Emergency Medicine, Ondokuz Mayıs University, Samsun, Turkey

***Corresponding author:** Erenler AK, Hıtit University Çorum Education and Research Hospital, Department of Emergency Medicine, Çorum, Turkey, Tel: + 222110001; Email: akemalerenler@hitit.edu.tr

Citation: Erenler AK, Ay MO and Baydın A (2016) Emergency Department Approach to the Growing Global Public Health Problem: Violence Against Women. Emerg Med Inves 2016; G115. DOI: 10.29011/2475-5605.000015

Received Date: 24 October, 2016; **Accepted Date:** 10 November, 2016; **Published Date:** 17 November, 2016

Abstract

Violence Against Women (VAW) is a growing problem threatens women's physical, mental and social well-being with a nature differing among cultures worldwide. In this review, we aimed to present prevalence and socio-cultural characteristics of VAW in different countries and make suggestions for Emergency approach.

Keywords: Violence against women; Abuse; Emergency Department

Introduction

Violence Against Women (VAW) is a public health problem, and at the same time, human rights violation all over the world[1-3]. The term 'VAW' involves the infliction or threatened infliction of physical, sexual and psychological/emotional violence, i.e. any type of aggression committed by a person with the intent of inflicting harm or exercising power and control over a woman or girl [4]. Common causes and types of VAW are listed in the table. Violence against women seems to be more often than predicted. According to UN reports, the prevalence of domestic violence in Belgium, the USA, Norway, New Zealand, South Korea, Colombia, and Guinea is 25%, 28%, 17%, 38%, 20%, 58%, and 67%, respectively [5]. In another study by the World Health Organization with 24,000 women in 10 countries (Bangladesh, Brazil, Ethiopia, Japan, Namibia, Peru, Samoa, Serbia and Montenegro, Thailand, and the United Republic of Tanzania) showed that 15%-71% of the participants were subjected to violence by their husband [6].

In this review, our aim is to identify socio-economical and cultural components of VAW and make suggestions to contribute to appropriate approach in the Emergency Department (ED).

Material and Methods

Scientific databases such as PubMed, Science Direct, Web of Science, and Google Scholar were investigated entering the word "violence against women" and country names from the con-

tinents. The search was performed by giving priority to recent (last 3 years) studies in order to compose a review in the light of current literature. Studies regarding domestic violence and violence from a partner were also included into the study. While reviews and researches on VAW were included, case reports were excluded in the review.

Europe

Violence against women in Europe is not only pronounced in low-income European countries such as Lithuania, Estonia and Latvia, also pronounced in economically advanced countries such as Finland, Denmark, Spain, Sweden and the United Kingdom of Great Britain and Northern Ireland. These results reveal that low income and the absence of gender equality are clearly not the explanation for the high frequency of victim-blaming attitudes in other European countries [7]. In a study in a gynecological emergency department in Rome, it was determined that there has been little change in the number of cases seen during the 15 year period [8].

In a study, VAW was categorized into three parts as socio-demographical characteristics, well-being related characteristics and marriage related characteristics, and socio-demographical characteristics were found to be the main cause of VAW [9].

Violence against women is a common public health problem in Turkey, but women were not organized regarding bringing it to the public's attention until the 1980s [10].

As the refugee crisis has arisen in Europe, women forced to migrate from their homelands have become subjects to violence

during their journey and/or on arrival in a destination country [11].

In a study among medical students, it was determined that 11.5% of the women had been sexually assaulted. In the same study, importance of educational programmes was underlined [12].

In a study, women in Spanish universities were investigated in regards to VAW and it was reported that 62% of the students know of or have experienced situations of sexual abuse within the university institutions, but only 13% identify these situations in the first place [13].

In a study, as the most serious consequence of VAW, ratio of death due to violence was found to be 12.9% [14]. A study from Spain revealed characteristics of the intimate related violence victims as minor acts of physical violence towards young adult women, either married or single, with children, unemployed and with a low family income level [15]. This is an important finding since young abused women are known to be three times more likely to suffer psychological distress and have somatic complaints, and use medication [16].

In a study conducted by a questionnaire in Belgium, Iceland, Denmark, Estonia, Norway, and Sweden, it was reported that an overall lifetime prevalence of any abuse was reported by 34.8% of the pregnant women. In the same study, the ratio for physical abuse ranged between 9.7-30.8% [17].

Prostitution in some European countries is also a factor for exposure to violence and consequent elevated mental health concerns [18].

In a study in Romania, controlling behaviour or emotional, physical, or sexual abuse against women by the partner was expressed as having occurred at least once in more than half of the cases [19].

Africa

In Rwanda a study indicated that VAW is increasing by the years [20]. In another study in Kenya, majority of the perpetrators were found to be husbands [21].

According to 2012-2013 Mali Demographic and Health Survey, controlling behavior, childhood exposure to violence, and husband's use of alcohol were reported to be predictors of physical and sexual abuse [22].

In a study, it was reported that most of the HIV-infected, pregnant VAW victims were more likely to be unmarried, substance users, and to score above the threshold for psychological distress and depression in Cape Town, South Africa [23].

In Tanzania the lifetime exposure to violence was found to be 65 % among ever-married or ever-partnered women and the most frequent abuse method was emotional [24]. A study from

Ghana revealed that risk of domestic violence was 41% higher for women whose husbands ever experienced their father beating their mother. Other predictors of potential VAW were found to be place of residence and alcohol use [25].

In a study in Nigeria, it was reported that economic reasons were the most common cause of VAW and risk was lower for women with an educated partner [26]. It was reported that VAW during pregnancy is correlated with low birth weight in Southeast Ethiopia [27]. In a study, VAW was found to effect younger women more frequently. The study also revealed the effect of cultural differences on VAW. Moslem women and those who are involved in traditional beliefs were more likely to expose VAW. Additionally women with low income were more likely to face violence [28]. In their study, Fidan et al. emphasize the importance of gender equality for VAW prevention [29].

America

Prevalence of domestic violence is increasing in the USA [30]. A study in Brazil revealed that most of the VAW were emotional type and VAW resulted in sexual intercourse for fear, depressive-anxious mood and drug use. Additionally, majority of the patients described themselves as black or mixed-race, having less than eight years' schooling, not being in work, having a partner at the time of the study or in the last 12 months, having children, or following a religious practice. In the study, it was reported that 31.3% of the women used of alcohol and one or more unlawful drugs, such as marijuana, crack or cocaine [31]. In a study by Cavanaugh et al., it was reported that VAW was related to sexual violence and high rates of recent sexual violence in this population and foreign-born Latina were more likely to experience violence when compared with those born in the USA [32].

Violence against women is also an important cause of economic burden in the USA [33]. Search for strategic plans emphasizing primary prevention, advancing the science of prevention, translating science into effective programmes still continue in the US [34].

Australia

In an Australian study, VAW was found to be strongly related to domestic relocation [35]. It was reported that Immigrant and refugee women who have resettled in Australia are known to face barriers accessing services aimed at preventing and responding to family violence [36]. In a study, it was reported that greater investment in violence reduction strategies is required and economically justified, particularly in young women. While anxiety and depression was reported to represent the greatest proportion of the disease burden attributable to violence, women subjected to violence tended to have other behavioural risk factors for chronic disease and the impacts extend to the whole family, in particular children

who can also suffer health impacts from witnessing violence [37]. In another study, 90% of participants had some education or training about intimate partner violence. Participating midwives generally reported a high level of knowledge about intimate partner violence but held misconceptions about risks and characteristics of perpetrators of violence [38]. Social contact and interaction are particularly appreciated by victims of VAW, as is instrumental support such as financial help, housing, and child care [39]. When speaking of measures in Australia, national bipartisan policies at federal, state and often local levels, an emphasis on prevention and early intervention in health services; innovative interventions to improve the responses of health professionals and the health system responses to violence are required. It continues to be difficult to achieve sustained changes or indeed to support a majority of health care professionals to identify and support women experiencing violence, their children and families [40].

Asia

In Iran psychological, physical, life threatening and sexual violence were found to be 59.7%, 33.2%, 10% and 39.3%, respectively. Significant difference was found between violence and some socio demographic characteristics including: age, years of marriage, occupation, education, smoking, number of children, satisfaction with baby sex and socio economic status [41]. In another study from Iran, the majority of the victims reported experiencing domestic violence and emotional violence was more prevalent than other kinds of violence. Additionally, lower education level, marriage at a younger age, shorter duration of marriage, fewer children, being a housewife, and husband's unemployment had a significant relationship with domestic violence against women [42].

Hajnasiri et al. investigated all the relevant articles on domestic violence in Iran published between the years 2000 and 2015 systematically. Based on the 31 articles, which represent a sample size of 15,514 persons, they estimated the prevalence of domestic violence to be 66% [43].

In Lebanon, Syrian refugee women's health, and specifically their sexual and reproductive health, was found to be severely affected. An increase in gender-based violence and early marriage, a lack of access to emergency obstetric care, limited access to contraception, forced cesarean sections, and high cost of healthcare services, all contribute to poor sexual and reproductive health resulting in violence [44]. The systematic use of sexual violence as a tactic of war is well-documented. Emergent narratives from the Middle East also highlight increasing risk and incidence of violence among displaced populations in refugee camps in countries bordering states affected by conflict [45].

Speaking of violence against women; Female Genital Mutilation/Cutting (FGM/C) may present as cutting, pricking, remov-

ing and sometimes sewing up external female genitalia for non-medical reasons. The practice of FGM/C is highly concentrated in a band of African countries from the Atlantic coast to the Horn of Africa, in areas of the Middle East such as Iraq and Yemen, and in some countries in Asia like Indonesia [46].

In a study in Japan, characteristics of sexual assailants were investigated and common features were found to be Blaming the Victim, Minimization, and Avoidance of Responsibility [47].

In Malaysia The prevalence of any form of violence against pregnant women was 35.9%, consisting of: any psychological (29.8%); any physical (12.9%); and any sexual (9.8%) violence. Violence against pregnant women was significantly associated with: women's use of drugs, having had exposure to violence during childhood, having a violence-supporting attitude, having two or more children; and having partners who were smokers, alcohol drinkers, or had controlling behavior [48]. In Vietnam significant risk was found to be associated with husbands' behaviour that supports male power (extra-marital relationships; fighting with other men) and alcohol use [49].

In a study in Saudi Arabia, it was reported that there was no significant correlation between socio-demographic characteristics and being abused or not. However, by further analysis we found more sexual abuse among non-working women [50].

A study in China revealed that a husband's gender traditionalism was positively associated with husband-to-wife physical assault only when the husband was coupled with a wife who has non-traditional gender attitudes [51].

In Hong Kong, sexual violence (13.4%) was more frequent than physical violence (11.7%). Women were more likely to be abused by men they knew (13.5%) than by strangers (8%) [52].

A study was conducted to investigate the association between Intimate Partner Violence against women and child morbidity in South Asia. The children of an abused mother were found to be more likely to experience diseases like fever, diarrhea and acute respiratory infection compared to the children of non-abused mothers [53]. A review of data from 81 countries revealed that South Asia has the second highest prevalence of violence (41.7%) [54].

In Sri Lanka, there appears to be a positive change in the perception of inter-personal violence, which is no longer viewed as an entirely private matter, encouraging more women to seek outside services. However, in moving forward, there is a need to address many of the barriers that prevent women from seeking legal redress and recourse from violence [55].

In Afghanistan, less than half reported abuse to health care providers or were asked by health care providers about the context of their injuries [56].

A study in Saudi Arabia revealed that the overall prevalence of domestic violence was 80.7 and 100.0% for physical and psychological violence, respectively. On studying the reasons for physical violence, half (50%) of the participants reported no clear cause, 19.2% reported failure to adequately care for children (such as cleaning, feeding, and dressing), and 7.8% reported causes related to poor scholastic achievement and couple conflict about appropriate approaches of upbringing of children. Suspicion on wife's fidelity was the most common form of psychological violence (21%) [57].

Emergency Approach and Conclusion

Violence against women is a complex issue involving social, economic and traditional components. Early identification of violence in families and the application of intervention programs by health and social services centers may prevent many of the undesired consequences of this VAW. There is additionally a critical need to determine and deal with the factors contributing to the emergence of VAW [42]. Etiological factors and characteristics of VAW are summarized in table 1.

Common causes of Violence against women
Lack of adequate legal sanctions
Social status overlooking violence
Inequality among genders
Unprotected human rights
Weak institutional community response
High criminal violence level
High armed conflict
Place of residence
Marriage-related factors
Alcohol/drug use
Childhood exposure to violence
Traditions
Domestic relocation
Lower education level
Marriage in a young age
Unemployment
Supported male power
Common types of Violence against women
Verbal abuse
Physical abuse
Sexual abuse
Emotional abuse
Behavior controlling
Female genital cutting

Table 1: Etiology and types of Violence against women.

In terms of the Middle East, democratization and women's rights can be traced through the recent events. Health service providers, particularly those who work in EDs should be aware of the impact of intimate violence women and be aware of women at

risk using the knowledge of maternal characteristics statistically associated with VAW. Detecting violence and screening for post-traumatic stress disorder in clinical settings like EDs may be beneficial for those who suffer from VAW. Assessment by specialists in forensic documentation and interpretation of injuries with the provision of balanced expert opinions for court purposes can result in a number of benefits for the victims and the criminal justice system, including an increase in the rate of successful prosecutions [58,59].

It is also important to improve gender equality through public education on gender relationships, educate women and provide economic opportunities, and eliminate common laws that sustain gender inequality as necessary steps to fight against VAW [29].

The importance of adopting standard violence concepts and promoting the recognition and evaluation of this form of violence amongst health care professionals, criminal investigators and forensic personnel must be emphasized [15,40]. Emergency departments may be the first places where victims contact an official.

Academics and politicians face many challenges in their efforts to reduce VAW. Investing in public education and awareness initiatives and constant monitoring are essential to assess the effectiveness of efforts to change public attitudes for victims VAW [60]. Suggestions to eliminate VAW are listed in Table 2.

To take measures for early detection of VAW
Screening for post-traumatic stress disorder following VAW
Improving gender equality
Increasing economical and educational status of women
Improving laws for VAW
Assessment of the efforts to change public attitude

VAW: Violence against women

Table 2: Suggestions to avoid violence against women.

Most of the women are known to visit EDs several times before the violent episode, often with traumas. Emergency Department medical and nursing staff should be prepared and trained to successfully identify and manage victims of violence.

References

1. Krantz G and Garcia-Moreno C (2005) Violence against women. *J Epidemiol Community Health* 59: 818-21.
2. Martín-Baena D, Montero-Piñar I, Escribà-Agüir V, Vives-Cases C (2015) Violence against young women attending primary care services in Spain: prevalence and health consequences. *Fam Pract* 32: 381-386.
3. Guedes A, Bott S, Garcia-Moreno C, Colombini M (2015) Bridging the gaps: a global review of intersections of violence against women and violence against children. *Glob Health Action* 9: 31516.
4. Moracco K, Runyan C, Bowling M, Earp JAL (2007) Women's experi-

ence violence: a national study. *Women's Health Issues* 17: 3-12.

- 5. WHO (2002) Multi-country study on women's health and domestic violence against women. Geneva: Department of gender and women's health, family and community health.
- 6. Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Watts CH, et al. (2006) Prevalence of intimate partner violence: findings from the WHO multicountry study on women's health and domestic violence. *Lancet* 368: 1260-1269.
- 7. Lokhmatkina NV, Agnew-Davies R, Costelloe C, Kuznetsova OY, Nikolskaya IM, et al. (2015) Intimate partner violence and ways of coping with stress: cross-sectional survey of female patients in Russian general practice. *Fam Pract* 32: 141-146.
- 8. Aleandri V, Bertazzoni G, Vetrano G, Carlomagno L, Osborn JF (2015) Violence against women in Rome: An analysis of cases treated in the gynecology emergency department of the hospital Umberto I between 1999 and 2013. *Ann Ig* 27: 665-677.
- 9. Özcan NK, Günaydin S, Çitil ET, et al. (2016) Domestic Violence Against Women In Turkey: A Systematic Review And Meta Analysis. *Arch Psychiatr Nurs* 30: 620-629.
- 10. Savran G. 80'li Yılların Kampanyaları ve Özel Alanın Politikası, Özgürliği Ararken: Kadın Hareketlerinde Mücadele Deneyimleri İçinde (Campaigns of 80's and politics of specialists, when we search for freedom: Experiences on struggles of women).
- 11. Freedman J (2016) Sexual and gender-based violence against refugee women: a hidden aspect of the refugee "crisis". *Reprod Health Matters* 24: 18-26.
- 12. Lüleci NE, Kaya E, Aslan E, Şenkal ES, Çiçek ZN (2016) Marmara University Medical Students' Perception on Sexual Violence against Women and Induced Abortion in Turkey. *Balkan Med J* 33: 173-180.
- 13. Valls R, Puigvert L, Melgar P, Garcia-Yeste C (2016) Breaking the Silence at Spanish Universities: Findings From the First Study of Violence Against Women on Campuses in Spain. *Violence Against Women* 22: 1519-1539.
- 14. Unal EO, Koc S, Unal V, Akcan R, Javan GT, et al. (2016) Violence against women: A series of autopsy studies from Istanbul, Turkey. *J Forensic Leg Med* 40: 42-46.
- 15. Regueira-Díéguez A, Pérez-Rivas N, Muñoz-Barús JI, Vázquez-Portoméne F, Rodríguez-Calvo (2015) MS4. Intimate partner violence against women in Spain: A medico-legal and criminological study. *J Forensic Leg Med* 34: 119-126.
- 16. Martín-Baena D, Montero-Piñar I, Escribà-Agüir V, Vives-Cases C (2015) Violence against young women attending primary care services in Spain: prevalence and health consequences. *Fam Pract* 32: 381-386.
- 17. Lukasse M, Schroll AM, Ryding EL, Campbell J, Karro H, et al. (2014) Prevalence of emotional, physical and sexual abuse among pregnant women in six European countries. *Acta Obstet Gynecol Scand* 93: 669-677.
- 18. Krumrei-Mancuso EJ (2016) Sex Work and Mental Health: A Study of Women in the Netherlands. *Arch Sex Behav*.
- 19. Rada C (2014) Violence against women by male partners and against children within the family: prevalence, associated factors, and inter-generational transmission in Romania, a cross-sectional study. *BMC Public Health* 14: 129.
- 20. Thomson DR, Bah AB, Rubanzana WG, Mutesa L (2015) Correlates of intimate partner violence against women during a time of rapid social transition in Rwanda: analysis of the 2005 and 2010 demographic and health surveys. *BMC Womens Health* 15: 96.
- 21. Onu CC, Dworkin SL, Ongeri LG, Oyaro P, Neylan TC, et al. (2016) Sexual violence against HIV-positive women in the Nyanza region of Kenya: Is condom negotiation an instigator? *J Acquir Immune Defic Syndr*.
- 22. Hayes BE and van Baak C (2016) Risk Factors of Physical and Sexual Abuse for Women in Mali: Findings From a Nationally Representative Sample. *Violence Against Women*: 1077801216658979.
- 23. Bernstein M, Phillips T, Zerbe A, McIntyre JA, Brittain K, et al. (2016) Intimate partner violence experienced by HIV-infected pregnant women in South Africa: a cross-sectional study. *BMJ Open* 6: e011999.
- 24. Kazaura MR, Ezekiel MJ, Chitama D (2016) Magnitude and factors associated with intimate partner violence in mainland Tanzania. *BMC Public Health* 16: 494.
- 25. Owusu Adjah ES and Agbemafie I (2016) Determinants of domestic violence against women in Ghana. *BMC Public Health* 16: 368.
- 26. Ansari U, Cobham B, Etim EM, Ahamad HM, Owan NO, et al. (2016) Insights Into Intimate Partner Violence in Pregnancy: Findings From a Cross-Sectional Study in Two States in Nigeria. *Violence Against Women*: 1077801216644072.
- 27. Demelash H, Nigatu D, Gashaw K (2015) A Case-Control Study on Intimate Partner Violence during Pregnancy and Low Birth Weight, Southeast Ethiopia. *Obstet Gynecol Int*: 394875.
- 28. Doku DT and Asante KO (2015) Women's approval of domestic physical violence against wives: analysis of the Ghana demographic and health survey. *BMC Womens Health* 15: 120.
- 29. Fidan A and Bui HN (2016) Intimate Partner Violence Against Women in Zimbabwe. *Violence Against Women* 22: 1075-1096.
- 30. Haqqi S and Faizi A (2010) Prevalence of domestic violence and associated depression in married women at a tertiary care hospital in karachi. *Procedia-Social Behav Sci*: 1090-1097.
- 31. de Barros ÉN, Silva MA, Falbo Neto GH, Lucena SG, Ponzo L, et al. (2016) Prevalence and factors associated with intimate partner violence among women in Recife/Pernambuco, Brazil. *Cien Saude Colet* 21: 591-598.
- 32. Cavanaugh CE, Messing JT, Amanor-Boadu Y, O'Sullivan CO, Webster D, et al. (2014) Intimate partner sexual violence: a comparison of foreign- versus US-born physically abused Latinas. *J Urban Health* 91: 122-135.
- 33. Max W, Rice DP, Finkelstein E, Bardwell RA, Leadbetter S (2014) The economic toll of intimate partner violence against women in the United States. *Violence Vict* 19: 259-272.
- 34. Graffunder CM, Noonan RK, Cox P, Wheaton J (2004) Through a public health lens. Preventing violence against women: an update from the US Centers for Disease Control and Prevention. *J Womens Health (Larchmt)* 13: 5-16.
- 35. Dillon G, Hussain R, Kibele E, Rahman S, Loxton D (2016) Influence of Intimate Partner Violence on Domestic Relocation in Metropolitan and Non-Metropolitan Young Australian Women. *Violence Against Women* 22: 1597-1620.
- 36. Vaughan C, Murdolo A, Murray L, Davis E, Chen J, et al. (2015) ASPIRE: A multi-site community-based participatory research project to increase understanding of the dynamics of violence against immigrant

and refugee women in Australia. *BMC Public Health* 15: 1283.

37. Cadilhac DA, Sheppard L, Cumming TB, Thayabaranathan T, Pearce DC, et al. (2015) The health and economic benefits of reducing intimate partner violence: an Australian example. *BMC Public Health* 15: 625.

38. Baird KM, Saito AS, Eustace J, Creedy DK (2015) An exploration of Australian midwives' knowledge of intimate partner violence against women during pregnancy. *Women Birth* 28: 215-220.

39. Tacket A, O'Doherty L, Valpied J, Hegarty K (2014) What do Australian Women Experiencing Intimate Partner Abuse Want From Family and Friends? *Qual Health Res* 24: 983-996.

40. Signorelli MC, Taft A, Pereira PP (2012) Intimate partner violence against women and healthcare in Australia: charting the scene. *Cien Saude Colet* 17: 1037-1048.

41. Abbaspoor Z and Momtazpour M (2016) Domestic Violence and Its Related Factors Based a Prevalence Study in Iran. *Glob J Health Sci* 8: 55206.

42. Mohamadian F, Hashemian A, Bagheri M, Direkvand-Moghadam A (2016) Prevalence and Risk Factors of Domestic Violence against Iranian Women: A Cross-Sectional Study. *Korean J Fam Med* 37: 253-258.

43. Hajnasiri H, Ghanei Gheshlagh R, Sayehmiri K, Moafi F, Farajzadeh M (2016) Domestic Violence Among Iranian Women: A Systematic Review and Meta-Analysis. *Iran Red Crescent Med J* 18: e34971.

44. Yasmine R and Moughalian C (2016) Systemic violence against Syrian refugee women and the myth of effective intrapersonal interventions. *Reprod Health Matters* 24: 27-35.

45. Alsaba K and Kapilashrami A (2016) Understanding women's experience of violence and the political economy of gender in conflict: the case of Syria. *Reprod Health Matters* 24: 5-17.

46. Muteshi JK, Miller S, Belizán JM3 (2016) The ongoing violence against women: Female Genital Mutilation/Cutting. *Reprod Health* 13: 44.

47. Hazama K and Katsuta S (2016) Cognitive Distortions Among Sexual Offenders Against Women in Japan. *J Interpers Violence*. pii: 0886260516669544.

48. Khaironisak H, Zaridah S, Hasanain FG, Zaleha MI (2016) Prevalence, risk factors, and complications of violence against pregnant women in a hospital in Peninsular Malaysia. *Women Health* 11: 1-23.

49. Jansen HA, Nguyen TV, Hoang TA (2016) Exploring risk factors associated with intimate partner violence in Vietnam: results from a cross-sectional national survey. *Int J Public Health*.

50. Al Dosary AH (2016) Health Impact of Domestic Violence against Saudi Women: Cross Sectional Study. *Int J Health Sci (Qassim)* 10: 165-173.

51. Cheung AK and Choi SY (2016) Non-Traditional Wives With Traditional Husbands: Gender Ideology and Husband-to-Wife Physical Violence in Chinese Society. *Violence Against Women* pii: 1077801216632615.

52. Bouhours B and Broadhurst R (2015) Violence Against Women in Hong Kong: Results of the International Violence Against Women Survey. *Violence Against Women* 21: 1311-1329.

53. Ferdousy EZ and Matin MA (2015) Association between intimate partner violence and child morbidity in South Asia. *J Health Popul Nutr* 33: 16.

54. Devries KM, Mak JYT, García-Moreno C, Petzold M, Child JC, et al. (2013) Global Health. The Global Prevalence of Intimate Partner Violence Against Women. *Science* 340: 1527-1528.

55. Guruge S, Jayasuriya-Illesinghe V, Gunawardena N, Perera J et al. (2015) Intimate partner violence in Sri Lanka: a scoping review. *Ceylon Med J* 60: 133-138.

56. Stokes S, Seritan AL, Miller E (2016) Care Seeking Patterns Among Women Who Have Experienced Gender-Based Violence in Afghanistan. *Violence Against Women* 22: 817-831.

57. Abo-Elfetoh NM and Abd El-Mawgod MM (2015) Violence against wives: a silent suffering in northern community. *J Egypt Public Health Assoc* 90: 87-93.

58. Fulu E and Miedema S (2015) Violence Against Women: Globalizing the Integrated Ecological Model. *Violence Against Women* 21: 1431-1455.

59. Nittis M, Hughes R, Gray C, Ashton M (2013) Domestic violence documentation project 2012. *J Forensic Leg Med* 20: 683-689.

60. Gracia E (2014) Intimate partner violence against women and victim-blaming attitudes among Europeans. *Bull World Health Organ* 92: 380-381.