

Research Article

Evaluation of Protein Levels in Diets of Salema porgy (*Sarpa salpa*) Juveniles

Merve Sahinyilmaz¹ and Murat Yigit^{2*}

¹Canakkale Onsekiz Mart University, Graduate School of Natural and Applied Sciences, Department of Aquaculture, Canakkale - Turkey

²Canakkale Onsekiz Mart University, Faculty of Marine Sciences and Technology, Departments of Aquaculture and Marine Technology, Canakkale - Turkey

*Corresponding author: Yigit M, Canakkale Onsekiz Mart University, Faculty of Marine Science and Technology, Terzioglu Campus. Canakkale-17100, Turkey. Tel: +905543132513, Fax: +902862180543, Email: muratyigit@comu.edu.tr

Citation: Sahinyilmaz M and Yigit M (2017) Evaluation of Protein Levels in Diets of Salema porgy (*Sarpa salpa*) Juveniles. J Fish Aquac Dev: JFAD-108. DOI:10.29011/JFAD-108/100008

Received Date: 27 February, 2017; **Accepted Date:** 28 March, 2017; **Published Date:** 5 April, 2017

Abstract

In the present study, the effects of different dietary protein levels on Salema porgy, *Sarpa salpa* [1] juveniles were investigated. Six iso-caloric (20 kJ g⁻¹ diet) diets with increasing protein levels (30, 37, 40, 47, 50, and 57%) were formulated. Each test diet was randomly fed to triplicate groups of 13 juvenile fish (initial mean weight 19.28±0.13 g) to satiety over 90 days. Growth performance and feed utilization were best with low dietary protein levels of 30 and 37%, but decreased with diets containing protein levels over 40%. Ammonia nitrogen excretion showed an increasing trend as dietary protein levels gradually increased, whereas retention rates of ammonia nitrogen per intake were highest in the low protein groups of 30 or 37%. The analyses of specific growth rate by broken-line regression indicated that the optimal dietary level of protein for salema porgy juvenile were 33.5% under the conditions applied in this study. As a result, *S. salpa* demonstrated better growth with low protein diets, showing that this marine fish could be a promising candidate for a sustainable and environment friendly aquaculture industry.

Keywords: Growth Performance; Feed Efficiency; Nitrogen Retention; Protein Requirement; Salema; *Sarpa salpa*

Introduction

The growing trend of marine aquaculture in southern European seas has doubled its production in the last ten years and reached about 276.000 tons with a total income of 1.783.000.000 US dollars in year 2014 [2]. It is estimated that the world population might reach 8.5 billion in year 2030 [3], where the need for human food will increase drastically. The aquaculture industry, with its increasing trend seems to be capable to supply an important amount of the food demand for human consumption. Nevertheless, seabream and seabass are the two main fish species in the Mediterranean with a production around 200.000 tons in Greece and Turkey [2], and the sales value of these two species are in pressure due to the high production rate and limited product diversity in the market. The introduction of new fish species in the market may trigger the demand and expand product diversity with new market opportunities. Salema porgy (*S. salpa*) is a member of the Sparidae family, known as a herbivorous fish species, feeding on plants, dis-

tributed around seagrass such as *Posedonia sp.* or *Cymodocea sp.* near the shore on sandy, or rocky sea bottom [4,5]. Salema porgy can be found in a wide range of area from shallow waters to 70 m deep water layers in the eastern part of the Atlantic (from the North Sea to Cape of Good Hope, the Canaries and Cape Verde Islands, in the Mediterranean and the Black sea), and in the western part of the Indian Ocean (from Mozambique to Cape of Good Hope) [6,7]. Schooling behavior of salema porgy around cage farms in the Mediterranean has been reported by [8], feeding on uneaten pellets that disperse from the fish pens, which is an indication that salema porgy can easily adapt to artificial pellet diets. From this point of view, salema porgy might be potential marine fish species for the Mediterranean aquaculture industry.

In contrast to gilthead seabream, or other sparid fishes, lower dietary protein requirements of salema porgy could be expected due to its herbivorous nature. Considering the rapid expansion of the world aquaculture industry and the disorganized or irregular state of the global capture fisheries which supplies the ingredients for aqua-diets, the demand for fishmeal and fish oil is likely to increase significantly. During the 2010-2030 period prices are ex-

pected to increase by 90 % for fishmeal and 70 % for fish oil according to [9]. For a sustainable development of the aquaculture industry, a gradual decline of capture fisheries as protein supply for fish feed production has been reported as essential [10]. Based on the increase of global fishmeal costs, [11] reported that fish meal usage in aqua-diets will decrease in the long term. Due, nowadays researchers have intensified their studies on the replacement of fishmeal or fish oil by less expensive alternative sources [12]. Besides, [9] suggested that in the face of higher fishmeal and fish oil prices, the substitution of fish species with less fishmeal requirements should also be considered and preferred for the marine aquaculture industry. Salema porgy (*S. salpa*) is a sparid fish, frequently seen around aquaculture cage farms in the Mediterranean and the Aegean Seas. [13] reported that, salema together with Striped mullet (*Mugil cephalus*) and white trevally (*Pseudocaranx dentex*) captured around fish cages in the Mediterranean had a stomach with pellets in great quantities. Furthermore, [8] also reported that salema schools around floating cage farms in the Mediterranean, and feeding on uneaten pellets. These observations strengthens the potential of salema porgy as a candidate marine fish species for the Mediterranean aquaculture industry.

Several reports on the ecology, reproductive biology, age-growth variation or geographic distribution of the wild populations of salema porgy [4,14-17] are available, however, to our knowledge so far, information on their nutritional requirements relative to their feeding habits is still lacking. Hence, this is the first attempt to assess the protein requirements and fed utilization of salema porgy with reference to growth performance, fish body bio-chemical composition and nitrogen budget under controlled culture conditions.

Materials and Methods

Experimental fish and rearing conditions

The feeding trial was conducted at the marine aquaculture research and development facilities of Marine Science and Tech-

nology Faculty at Canakkale Onsekiz Mart University (Dardanos-Canakkale, Turkey). Initial and final fish were weighed individually (precision 0.01 g). At intervals of 30 and 60 days during the course of the feeding trial however, fish were mass weighed in buckets filled with seawater in order to avoid handling and netting stress. Before weighing, fish were deprived of feed for one day. Experimental fish with initial mean weight of 19.28 ± 0.13 g were placed into 18 circular polyethylene tanks with a water volume of 200 L. A factorial design of 6x3 was applied and a total of 234 fish were randomly stocked in six groups of tanks with 13 fish per tank, and 3 replicates per treatment. Experimental fish were adapted for a period of 1 month to the culture conditions prior to start of feeding trial, which was initiated when all fish accepted pellets. Seawater was supplied to the tanks at a flow rate of 28 L min^{-1} . Aeration was continuously supplied by air-stones and the photoperiod regime was a natural light course ($40^{\circ}04'37.47''\text{N } 26^{\circ}21'39.04''\text{E}$). Throughout the feeding trial, ambient water parameters such as temperature, salinity, dissolved oxygen, pH were measured periodically using a YSI multi-probe water analyser. Total ammoni-anitrogen ($\text{NH}_3\text{-N}$) was determined by the Nessler method using a HANNA (HI 2221) /portable spectrophotometer (HANNA Instruments Co., Padova, Italy).

Experimental Diets and Feeding

Practical diets were formulated with commercially available ingredients and produced at the laboratories of Canakkale Onsekiz Mart University, Faculty of Marine Science and Technology in Canakkale, Turkey. All the test diets were formulated to be isocaloric on a gross energy (20.0 kJ/g diet) basis and to contain increasing levels of protein (30, 37, 40, 47, 50 and 57 %). Total n-3 Highly Unsaturated Fatty Acid (HUFA) contents averaged 3.6 g/kg for all test diets. Brown fish meal (anchovy, Blacksea origin) was used as a sole protein source. Ingredients and chemical composition of test diets are given in (Table 1), and the amino acid profiles of the experimental diets are presented in (Table 2).

Experimental diet / Protein level						
Ingredient (g/kg DM)	D1/30	D2/37	D3/40	D4/47	D5/50	D6/57
Fish meal ¹	410	490	565	647	730	810
Corn starch	50	50	50	50	50	50
Dextrin	405	335	265	190	115	43
Fish oil (FO)	90	80	75	68	60	52
Vit-min mix ²	40	40	40	40	40	40
Cholin chloride	5	5	5	5	5	5
Total	1000	1000	1000	1000	1000	1000
Proximate composition (% DM, dry matter)						
Dry matter	91.6	91.0	91.3	91.1	91.6	91.5
Crude protein	29.4	36.7	41.4	46.2	52.1	57.2
Crude lipid	16.0	17.9	17.1	19.3	18.4	19.7
Crude ash	4.46	5.07	6.67	7.53	8.64	9.14
NFE ³	38.8	28.3	23.1	15	9.46	2.41
GE (kJ/g diet) ⁴	19.6	20.2	20.1	20.7	20.8	21.2
GE (kcal/g diet)	4.68	4.83	4.8	4.94	4.96	5.08
P:E (mg/kJ) ⁵	15	18.2	20.6	22.3	25.1	26.9
PE/TE	0.35	0.43	0.49	0.53	0.59	0.64
Crude lipid in FM (%)	8.5	8.5	8.5	8.5	8.5	8.5
Lipid from FM (%)	3.49	4.17	4.8	5.5	6.21	6.89
Σ FO in diet (%)	12.5	12.2	12.3	12.3	12.2	12.1
n-3 HUFA in FO (%) ^a	29.8	29.8	29.8	29.8	29.8	29.8
Σ n-3 HUFA in diet (%)	3.72	3.62	3.66	3.66	3.63	3.6

¹Anchovy meal, Blacksea-Turkey
²Vitamin mixture (per 1 mg): Vit.A 65.000 IU, Vit.D3 45.000 IU, Vit.E 25 IU; Vit.K3 5 mg, Vit.B1 12.5 mg, Vit.B2 12.5 mg, Vit.B6 15 mg, Vit.B12 0.025 mg and ascorbic acid 120 mg; Mineral mixture (per 1 mg): Ca 100 mg, P 50 mg, K 30 mg, Na 20 mg, Mg 10 mg, Fe 22 mg, Zn 3 mg, Mn 3 mg, Cu 1.8 mg, Co 0.15 mg, I 0.12 mg, Se 0.05 mg, DL-calcium pantothenate 40 mg, niacin 50 mg, folic acid 2.5 mg, biotin 0.08 mg and inositol 75 mg.
³Nitrogen free extract = 100 - (crude oil + crude ash + crude protein)
⁴Gross energy; calculated based on energy fuels of 23.6 kJ/g protein, 39.5 kJ/g lipid and 17.2 kJ/g NFE.
⁵Protein-enerji ratio = mg protein / kJ energy
⁶PE/TE = energy from protein / total energy
 Σ n-3 HUFA in diet (g/kg) = (Σ fish oil in diet, g/kg) x (% n-3 HUFA in fish oil used)

Table 1: Ingredients and proximate composition of the experimental diets.

Amino acid (%/dry matter)	Experimental diet /Protein level						
	Fish meal ^a	30	37	40	47	50	57
Arginine	4.11	1.69	2.01	2.32	2.66	3	3.33
Lysine	5.49	2.25	2.69	3.1	3.55	4.01	4.45
Histidine	1.76	0.72	0.86	0.99	1.14	1.28	1.43
Isoleucine	3.38	1.39	1.66	1.91	2.19	2.47	2.74
Leucine	5.43	2.23	2.66	3.07	3.51	3.96	4.4
Valine	3.81	1.56	1.87	2.15	2.47	2.78	3.09
Methionine	2.16	1.16	1.38	1.59	1.82	2.06	2.28

Phenylalanine+Tyrocine	5.47	2.24	2.68	3.09	3.54	3.99	4.43
Threonine	3	1.23	1.47	1.7	1.94	2.19	2.43
Tryptophan	0.82	0.34	0.4	0.46	0.53	0.6	0.66
ΣEAA	35.4	14.8	17.7	20.4	23.4	26.3	29.2
^a according to N/A = not available							

Table 2: Amino acid profiles of test diets with increasing levels of protein (g/16 g N).

Initially, all ingredients including oil were mixed with a food mixer for 20 min, then tap water was added in order to prepare a suitable pulp, that was made into a 2 mm sized pellets with a meat grinder. The pelleted diets were then dried to a moisture content of 80-90 g/kg at 40°C in a drying chamber. The test diets were then stored in a freezer (-25°C) until use. Experimental fish were hand fed until satiation twice a day at 09:00 and 16:30 hours for a total of 90 days. Special attention was given to be certain of the even distribution of pellets by all fish in the tanks, and feeding lasted for about 15-20 min. When fish refused feeding, it was accepted as a sign of satiety and feeding was stopped in order to avoid overfeeding. In all tanks, the feed intake was recorded daily by subtracting the feed distributed from the initial weight of feed.

Sampling and Analytical Methods

Prior to the start of the experiment, 10 fish from the initial pool were anesthetized in a high dose MS-222 (100 mg/L) and stored in polyethylene bags in a freezer (-25°C) for subsequent analysis. At the end of the experiment, the same protocol of sampling was followed for each tank. Five fish per tank (15 fish per treatment) were randomly withdrawn for comparative analysis of fish whole body (dry matter, protein, lipid, ash) and calculation of nutrient retention rates and nitrogen budget. All analyses were performed in triplicate and samples were prepared by homogenizing fish whole body in a kitchen blender. Chemical analyses of test diets and fish whole body were conducted according to [18] guidelines as follows: for dry matter, drying in an oven at 105°C for 24 h until constant weight were obtained; for protein (Nx6.25) by Kjeldahl method after acid digestion; for lipids by ethylether extraction in a Soxhlet System; for ash by incineration in a muffle furnace at 550°C for 12 h. The NFEs were calculated by subtracting the sum of protein, lipid and ash from hundred.

Statistical Analysis

The results were given as mean±Standard Deviation (SD) and differences of group means were compared by one-way ANOVA. Significance level of $p < 0.05$ was applied for all data. In order to figure out the optimum dietary protein level that matches with the maximum growth rate, a third order polynomial regression between dietary protein and growth rate values was applied [19].

Results and Discussion

At the end of the 90 days growth experiment, survival rates were over 85% for all treatment groups, indicating that dietary protein levels did not affect fish survival. Best growth performance in salema porgy juveniles were obtained when fed a diet with 37% protein. This was followed by the 30% and 40% diet groups, respectively. No significant difference ($p > 0.05$) was found between final body weight of fish fed the 37% protein diet and those fed diets with 30% or 40% protein levels. However, dietary protein levels over 40% significantly ($p < 0.05$) reduced the growth rates (Figure 1). The best Specific Growth Rates (SGR) were obtained in fish fed the 37% protein diet, which demonstrated significantly better ($p < 0.05$) performance compared to the higher protein diet. Even though there was no significant difference in SGRs between the 30% and 37% protein diets, the latter performed about 15% better than the 30% dietary protein group. A gradual decline was observed in percent feed intakes with the decrease in dietary protein levels. The highest feeding rate of 0.69% ($p < 0.05$) was recorded in fish fed diets containing 37% protein. Based on the polynomial regression analyses [19] used for the relation between dietary protein levels and the SGRs, it was recorded that the optimum protein requirement for juvenile salemaporgy was about 33.5% of the diet under the conditions applied in this study (Figure 2). The values for Protein Efficiency Rates (PER) followed the same trend, with higher rates ($p < 0.05$) for the best performing diet groups of 30% and 37%, which demonstrated significantly lower ($p < 0.05$) Feed Conversion Rates (FCR) compared to the higher protein diets (Table 3).




Figure 1: Growth trend of salemaporgy fed diets with six different protein levels for 90 days. Values with different letters are significantly different ($p < 0.05$).




Figure 2: Optimum dietary protein requirement of salema porgy (*Sarpa salpa*) juvenile by polynomial regression between dietary protein levels and specific growth rates. Values with the different letters are significantly different ($p < 0.05$). (Optimum protein level shown with arrow).

Experimentaldiet/protein level						
D1/30	D2/37	D3/40	D4/47	D5/50	D6/57	D5/50
IBW	19.3±0.08	19.2±0.04	19.2±0.04	19.3±0.10	19.2±0.07	19.5±0.14
FBW	28.9±1.03 ^b	30.5±1.12 ^b	27.0±1.45 ^{ab}	25.8±0.44 ^a	25.9±1.09 ^a	25.2±1.04 ^a
SGR	0.45±0.04 ^{bc}	0.51±0.04 ^c	0.38±0.06 ^{ab}	0.32±0.02 ^a	0.33±0.04 ^a	0.28±0.04 ^a
FI	0.59±0.02 ^c	0.69±0.02 ^d	0.61±0.04 ^c	0.51±0.02 ^a	0.55±0.01 ^b	0.52±0.03 ^{ab}
FCR	1.35±0.18 ^{ab}	1.24±0.12 ^a	1.66±0.35 ^{ab}	1.89±0.63 ^{ab}	1.71±0.26 ^b	1.88±0.37 ^b
PER	2.56±0.33 ^c	2.22±0.23 ^c	1.50±0.32 ^b	1.22±0.35 ^{ab}	1.14±0.18 ^{ab}	0.96±0.19 ^a
SR	87.18	92.31	89.74	84.62	84.62	87.18

* Values with different superscript letters in the same line are significantly different at $p < 0.05$ level.
 IBW: Initial Body Weight (g); FBW: Final Body Weight (g)
 SGR (Specific Growth Rate, % growth per day) = $((\ln W_2 - \ln W_1) / (t_2 - t_1)) \times 100$
 FI (Percent Feed Intake, % per day) = $(\text{total feed intake} / ((W_1 + W_2) / 2) / \text{day}) \times 100$
 FCR (Feed Conversion Rate) = $\text{feed intake (g)} / \text{weight gain (g)}$
 PER (Protein Efficiency Rate) = $(\text{weight gain (g)} / \text{protein intake (g)})$
 SR (Survival Rate, %) = $(\text{number of remaining fish} / \text{number of initial fish}) \times 100$

Table 3: Growth performance and feed utilization of salema porgy fed the experimental diets for 90 days (means ± SD)*.

During the first 2 month of the trial, growth of salema porgy was relatively low, however after the 60 days of the feeding trial, the growth showed an increasing trend compared to the initial performance. Since the culture conditions were the same throughout the study, the acceleration of growth performance in the second month with an increasing trend in the third month of the trial might be attributed to the week adaptation of salema juveniles in tank environment. Eventhough the experimental fish were adapted for a period of 1 month to the culture conditions, and the feeding trial initiated when all fish accepted pellets, it seems that salema juveniles might need a longer acclimatization period to tank conditions of certain sizes. For instance, the best performing group in the present study showed a SGR of 0.5 %/day throughout the feeding trial, while fish growth during the last period of 30-days resulted in an increased growth of 0.9 %/day. The accelerated increase of growth performance after the 60-days of the feeding trial might be an indication for a better growth performance of salema when a longer adaptation period were applied.

The maximum SGR (0.51 %/day) obtained for salema porgy in the present study was higher than an earlier report on axillary seabream (*Pagellus acarne*), an other candidate sparid fish for aquaculture (0.23 %/day, Yigit et al., 2016). Korkutand Balkı (2004) reported SGR variation of between 0.32 and 1.04 %/day for gilthead seabream under commercial cage farm conditions in the Aegean Sea. Similar to our findings in the present study for Salema porgy, [20,21] also reported other sparid candidates such as White seabream (*Diplodus sargus*) (0.89 %/day) and Zebra seabream (*Diplodus cervinus*) (0.8 %/day) as slow growing marine species, respectively. On the other hand, [21,22] recorded higher SGRs (1.22 %/day and 1.54 %/day) for two-banded seabream (*Diplodus vulgaris*) and sharpnout seabream (*Diplodus puntazzo*), respectively.

Physico-chemical water parameters recorded in the present study were comparable and within the acceptable limits reported by [23] for a recirculating aquaculture system (Table 4).

Parameter	Unit	Acceptable limits*	Present study
pH	-	6.5-7.5	7.55±0.6
Temperature	°C	Species specific	12.76±2.24
Salinity	ppm	Species specific	25.64±0.84
Oxygen	O ₂ , % (mg/L)	70-100	88 (7.98±1.3)
Ammonium	NH ₃ , mg/L	0-2.5 (pH influenced)	0.27±0.06
*Bregnballe, 2015			

Table 4: Acceptable limits for different physico-chemical water quality parameters in a recirculating system and recorded values in the present study.

Water temperature during the course of the present study (12.76±2.24°C) was relatively low compared to the data reported in the southern Aegean Sea where the average annual seawater temperature is in a range between 12.4-25°C [24]. Discrepancies between different reports might be due to rearing conditions such as water temperature, dissolved oxygen, salinity, fish stocking rates, feeding methods, or a combination of these factors [25,26], as well as to fish species and fish size [27].

Dietary protein requirements estimated based on growth performance (percent growth per day) or nitrogen retention (per N-intake) corresponded to 37% and 30%, respectively. Similar to our findings, protein requirements for two-banded seabream (*Diplodus vulgaris*) were reported between 35-36% in earlier studies [22,28], while a lower protein level of 27% was recorded for juvenile white seabream (*Diplodus sargus*) [20]. In contrast, higher protein levels for best growth were found for other sparids such as Blackfin seabream (*Acanthopagrus berda*) 42% [29], Zebra seabream (*D. cervinus*) 43.8% [21], and Sharpnose seabream (*Diplodus puntazzo*) 43-47% [21,30], respectively. [31] reported that generally fast growing fish species require higher protein levels in their diets compared to the slow growing species. Apart from this, carnivorous fish species such as Gilthead seabream (*S. aurata*), European seabass (*Dicentrarchus labrax*), rainbow trout (*Onchorhynchus mykiss*), Common dentex (*Dentex dentex*), Japanese flounder (*Paralichthy solivaceus*), or the red porgy (*Pagrus pagrus*) are reported to require higher levels of dietary protein (45-55%) [32-37]. The lower protein requirement of salema porgy

compared to other sparids or other marine fishes could be attributed to their herbivorous feeding nature, whereas the others requiring higher protein diets are either carnivorous or omnivorous species. This is in agreement with the statement of [38], indicating that carnivorous fishes have higher protein requirements compared to omnivorous or herbivorous species.

The regulation of feed intake in order to meet energy demand is general condition for fish [39]. This condition has also been reported by [20] for white seabream, by [21] in zebra seabream, [40] and [21] in sharpnose seabream, and [29] in Blackfin seabream. Our finding for the feed intake in salema porgy is in agreement with earlier reports, in terms of increased feed intake ($p < 0.05$) with the decrease of protein levels in the diets. In the present study, the FCRs of Salema porgy linearly increased with the declining dietary protein levels. This is in agreement with the findings of [20] in white seabream, [21] in Sharpnose seabream, and [29] in Blackfin seabream. In contrast, [21] and [41] reported increased FCRs in Zebra seabream and Gilthead seabream, when dietary protein levels also increased. The FCRs found in our study (1.24-1.89) were comparable with those reported earlier for two banded seabream (*D. vulgaris*) (1.50-1.80, 1.36-2.96, 1.67-1.92; [28] ab [42], respectively), in Zebra seabream (1.69-3.33) [21], and in Gilthead seabream (0.91-3.06, 1.14-3.73, 1.22-1.74, 1.24-1.48, 1.37-1.53 [25,26,41,43,44], respectively). Higher FCR of 2.51 was found in axillary seabream (*Pagellus acarne*), introduced as a new candidate species for the Mediterranean aquaculture by [45], while [46] reported lower FCRs of 1.1-1.2 in gilthead seabream. Similar to the growth performance recorded in the present study, FCRs and PERs followed the same trend with better results in fish fed lower protein diets. These findings in the present study were in agreement with earlier reports on European eel *Anguilla anguilla* [47], Yellow snapper *Lutjanus argentiventris* [48], and two-banded Seabream *D. vulgaris* [28] in terms of decreasing PERs with increased dietary protein levels.

The nitrogen retention rates per intake in fish fed diets with 30% and 37% protein were significantly ($p < 0.05$) higher than those fed the higher protein diets. In contrast, excretion rate of nitrogen per intake were lowest for the fish fed on lower protein diets of 30 and 37% and showed a significant increase when dietary protein levels rose over 40% (Table 5).

Experimental diet / protein level						
N balance (mg g ⁻¹ production)	D1/30	D2/37	D3/40	D4/47	D5/50	D6/57
NI	73.5±18.4 ^a	83.4±12.1 ^a	110.2±23.5 ^{ab}	119.8±12.7 ^b	142.2±21.4 ^{bc}	171.6±33.6 ^c
NR	30.1±2.54 ^a	30.3±1.83 ^a	31.2±5.81 ^a	31.1±2.47 ^a	27.9±2.97 ^a	30.0±5.16 ^a
TNE	43.4±17.3 ^a	53.0±13.9 ^a	79.0±17.7 ^{ab}	88.7±10.9 ^b	114.3±18.4 ^{bc}	141.6±28.6 ^c
NR (%NI)	42.4±8.7 ^d	37.1±7.12 ^d	28.4±0.90 ^c	26.0±1.73 ^c	19.7±0.99 ^b	17.6±0.81 ^a
TNE (%NI)	57.6±8.7 ^a	62.9±7.12 ^a	71.6±0.90 ^b	74.0±1.73 ^b	80.3±0.99 ^c	82.5±0.81 ^d
* Values with different superscript letters in the same line are significantly different at p<0.05 level. NI (Nitrogen Intake, mg/g production) = (protein intake / 6.25) / (W2 - W1) NR (Nitrogen Retention, mg/g production) = (total g protein retained in fish / 6.25) / (W2 - W1) TNE (Nitrogen Total Excretion, mg/g production) = (nitrogen intake(g)-nitrogen retention(g))/(W2-W1) NR (%NI, Nitrogen Retention as percent of nitrogen intake) = 100 x (N retention / N intake) TNE (%NI, Nitrogen Total Excretion as percent of nitrogen intake) = 100 x (N excretion / N intake) W2: Final Fish Weight (g), W1= Initial Fish Weight (g)						

Table 5: Nitrogen (N) balances of salema porgy fed to satiation the experimental diets during 90 days(means±SD)*

The excessive supplement of dietary animal proteins may result in increased nitrogen excretion. The incorporation of dietary animal protein or lipids at an optimum level may support the aquaculture industry economically and environmentally [22]. In the present study, dietary protein levels over 40 % resulted in a significant increase of nitrogen excretion, which can be explained by the elevated protein catabolism led to higher ammonia excretion rates in fish fed excessive dietary protein. This finding was also supported by the PERs in the present study with better protein utilization when fed diets lower than 40% protein. Our findings regarding nitrogen retention rates per intake (37-42 %) in best performing protein groups are in close agreement with earlier reports in European seabass fed different ration levels (36-43%; [49], in rainbow trout (18-46%; [50], Atlantic turbot (28-36%, [51]; 36-42%, [52], the Black Sea turbot (38-40%, [53]; 19-41%, [54]; 29-30%, [55]. Lower retention rate of nitrogen per intake have been reported in Blackfin seabream (20-40%, [29], zebra seabream (19-26%, [21], European seabass (23-32%, [56]; 16-26%, [57]. Reported that the optimal protein level in fish diet might be affected by the amino acid composition of the test proteins [58]. In earlier studies, it has been reported that feeding fish with diets over the requirement level may result in an increased protein catabolism [59], induced with the increase of hepatic activity of alanine aminotransferase, aspartate aminotransferase, and glutamate dehydrogenase enzyme activities [60-62]. In the present study, eventhough enzyme activities were not investigated, the reason for the higher nitrogen excretion rates in experimental fish fed higher levels of dietary protein might be attributed to the increased protein catabolism due to the excessive protein levels in the diets.

To our knowledge so far, there is no data available on the Essential Amino Acid (EAA) requirements of salema porgy. Considering the best performing diet of 37%, and the reduction in fish growth when fed in excess of requirements, might also be linked to an excessive dietary EAAs for the test diets containing protein lev-

els over 40 %. Hence, based on the findings in the present study, it might be assumed that the amino acid profile of the best performing diet (37%) is close to ideal EAA profile for salema porgy juveniles. Because, at this level of dietary protein, there were no limitation of amino acids in the test diets, that otherwise could have resulted in growth limitations of fish. Actually, fish diets below ideal protein profile lacking in one or more EAA can lead to reduced feed intake and growth performance, depress protein or amino acid retention, due to higher protein and amino acid catabolism, which in turn lead to increased nitrogen waste and deterioration of environment waters [21,29,50,59,63].

A slight decrease in fish whole body protein was observed when dietary protein levels increased over 40% level, however no significance (p>0.05) was recorded among the experimental groups. Crude lipid contents of fish body followed the same trend with no significant differences (p>0.05) among test groups. Fish body ash contents tended to increase with increasing levels of dietary protein, however these differences were not significantly (p>0.05) important as well (Table 6).

Diets	Hepatosomatic Index	Viscerasomatic Index
Initial	1.68±0.66	6.91±1.14
D1/30	1.52±0.27	6.11±0.52
D2/37	1.51±0.50	6.36±0.87
D3/40	1.53±0.21	6.37±0.41
D4/47	1.60±0.64	6.71±0.92
D5/50	1.83±0.39	7.27±0.67
D6/57	1.66±0.16	6.49±0.13
* Values with no superscript letters in the same row are not significantly different at p<0.05 level		

Table 7: Body morphological indices of salema porgy juveniles fed experimental diets with increasing protein levels for 90 days (means±SD)*.

Hepatosomatic Index (HSI) = (liver weight / total weight) x 100

Viscerasomatic Index (VSI) = (viscera weight / total weight) x 100

The liver is known to have a function as the deposition site for fat and glycogen in fish [64] and [65]. Reported that dietary carbohydrates stimulate glycolysis, glycogenesis and lipogenesis, while reducing protein catabolism and gluconeogenesis [66]. In the present study, protein levels and NFEs (soluble carbohydrate of the feed) of the test diets were negatively correlated, with increasing carbohydrates at decreasing levels of dietary protein. Due, the increased carbohydrate levels in our test diets with lower dietary protein might have stimulated the lower protein catabolism. It has been reported that HSI is positively correlated with dietary carbohydrate levels, while inversely related to dietary protein [21,67-69]. In the present study however, an adverse relation between HSI and carbohydrate level, but positive correlation with dietary protein levels were observed. The VSI showed similar trend as the HSI in the present study (Table 6).

Considering that the best growth was obtained with the low protein diets (30-40%), which were higher in NFEs (23-38% vs 2-15%) but lower in protein to energy (P:E) ratio (15-20 mg/kJ vs 22-27 mg/kJ), compared to the higher protein diets (47-57%) might be attributed to the herbivorous nature of salema porgy and also linked to a hypothesis that salema porgy might prefer low-protein but high-energy diets for a best growth performance, as also reported by [70] in two-banded seabream. However, the experimental diets in the present study were formulated with a single lipid level. Further studies are encouraged to assess dietary lipid and carbohydrate levels for salema porgy with experimentations at different water temperature regimes.

Conclusion

In the present study, optimum dietary protein requirement of salema porgy juveniles by polynomial regression between protein levels and growth rates was found as 33.5%, indicating that this low level of dietary protein is optimum for maximum growth and feed conversion ratio in salema porgy juveniles. Increasing the dietary protein over 40% seems to induce a decline on weight gain, and negatively affect the protein efficiency as well as nitrogen retention rates. As a marine fish species with low protein requirements, salema porgy might be a promising candidate for the Mediterranean aquaculture industry, with the less use of fishmeal based protein sources, that in long run might benefit the global aquaculture in terms of economically sustainable and environment friendly way.

Acknowledgement

CanakkaleOnsekiz Mart University, Scientific Research Projects Commission (COMU-BAP) is acknowledged for the support of laboratory equipments used in this study with the Project

ID: 256 and Project Code: FAY-2014-256. COMU, Faculty of Marine Science and Technology is acknowledged for the use of experimental facilities throughout the study, which has been performed in partial fulfillment of requirements for the degree of Master of Science of the first author.

References

1. Seabream (*Sparus aurata* Linnaeus, 1758) Iraq Aquaculture Journal 1: 25-35.
2. FAO (2016) Fisheries and Aquaculture Information and Statistics Branch, Food and Agriculture Organization of the United Nations, Global Production Statistics.
3. United Nations (2015) Probabilistic population projections based on the World population prospects: The 2015 Revision. United Nations, Population Division DESA.
4. Criscoli A, Colloca F, Carpentieri P, Belluscio A, Ardizzone G (2006) Observation on the reproductive cycle, age and growth of the salema, *Sarpa salpa* (Osteichthyes: Sparidae) along the western central coast of Italy. *Scientia Marina* 70: 131-138.
5. Steele L, Darnell KM, Cebrián J, Sanchez-Lizaso JL (2014) *Sarpa salpa* Herbivory on shallow reaches of *Posidonia oceanica* beds. *Animal Biodiversity and Conservation* 37: 49-57.
6. Smith MM, Heemstra PC (Eds.) (1986) *Smiths' Sea Fishes*. Berlin: Springer-Verlag 580-594.
7. Bauchot ML, Hureau JC, Quérou JC, Hureau JC, Karrer C, et al. (1990) Sparidae. In: (Eds.) Checklist of the fishes of the Eastern Tropical Atlantic Catalogue des poissons de l'Atlantique tropical oriental (Clofeta), vol 3. Unesco, Paris 790-812.
8. Colomi A, Diamant A, Woo PTK, Bruno DW (Eds.) (2014) Infectious diseases of warm water fish in marine and brackish waters. In: *Diseases and Disorders of Finfish in Cage Culture* (2nd ed.) Wallingford UK: CABI Publishing 155-192.
9. Msangi S, Kobayashi M, Batka M, Vannuccini S, Dey MM, et al. (2013) *Fish to 2030*. Washington (USA): Prospects for Fisheries and Aquaculture. (The World Bank Publication; report no. 83177-GLB).
10. Nengas I, Alexis MN, Davies SJ (1996) Partial substitution of fishmeal with soybean meal products and derivatives in diets for Gilthead seabream *Sparus aurata* (L.) *Aquaculture Research* 27: 147-156.
11. Tacon AGJ, Metian M (2008) Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285: 146-158.
12. Yigit M, Ergün S, Türker A, Harmantepe B, Erteken A (2010) Evaluation of soybean meal as a protein source and its effect on growth and nitrogen utilization of black sea turbot (*Psettima setacea*) juveniles. *Journal of Marine Science and Technology Taiwan* 18: 682-688.
13. Neofitou N (2016) Waste feed from fish farms of the Eastern Mediterranean and attraction of wild fish. *Universal Journal of Geosciences* 4: 112-115.
14. Villamil MM, Lorenzo JM, Pajelo JG, Ramos A, Coca J (2002) Aspects of the history of salema, *Sarpa salpa* (Pisces, Sparidae), off the Canarian Archipelago (central-east Atlantic) *Environmental Biology of Fishes* 63: 183-192.
15. Matic-Skoko S, Kraljević M, Dulčić J, Pallaoro A (2004) Growth of juvenile salema, *Sarpa salpa* (Teleostei: Sparidae), in the Kornati Archipelago, eastern Adriatic Sea. *Scientia Marina* 68: 411-417.

16. Dobroslavić T, Zlatović A, Bartulović V, Lučić D, Glamuzina B (2013) Diet overlap of juvenile salema (*Sarpa salpa*), bogue (*Boops boops*) and common two-banded sea bream (*Diplodus vulgaris*) in the south-eastern Adriatic. *Journal of Applied Ichthyology* 29: 181-185.
17. Bayhan B, Kara A (2015) Length-weight and length-length relationships of the salema *Sarpasalpa* (Linnaeus, 1758) in Izmir Bay (Aegean Sea of Turkey). *Pakistan Journal of Zoology* 47: 1141-1146.
18. AOAC (1998) Association of Official Analytical Chemists, Official Methods of Analysis of AOAC International: Maryland USA.
19. Brett JR, Grove TDD (1979) Physiological energetics in fish. In: Hoar WS, Randall DJ, Brett JR (Eds.) *Fish Physiology* Academic Press, New York: 279-352.
20. Sá R, Pousão-Ferreira P, Olive-Teles A (2008) Dietary protein requirement of White sea bream (*Diplodus sargus*) juveniles. *Aquaculture Nutrition* 14: 309-317.
21. Coutinho F, Peres H, Castro CO, Oliva-Teles A (2012) Dietary protein requirement and intermediary metabolism response to protein/carbohydrate ratio in zebra seabream (*Diplodus cervinus*, Lowe 1838) juveniles [Master of Science Thesis] Porto Portugal: Faculty of Science, University of Porto.
22. Ozório ROA, Valente LMP, Correia S, Pousao-Ferreira P, Damasceno-Oliveira A, et al. (2009) Protein requirement for maintenance and maximum growth of two-banded seabream (*Diplodus vulgaris*) juveniles. *Aquaculture Nutrition* 15: 85-93.
23. Bregnballe J (2015) A guide to recirculation aquaculture, an introduction to the new environmentally friendly and highly productive closed fish farming systems. Publication of Food and Agriculture Organization of the United Nations (FAO) and EUROFISH International Organisation.
24. Hossu B, Korkut AY, Salnur S (2005) Investigation on feeding tables for seabass (*Dicentrarchus labrax* L., 1758) in net-cage (Pinar Marine Company) culture. *Mediterranean Fish Nutrition*, in: Cahiers Options Méditerranéennes, edited by Montero D, Basurco B, Nengas I, Alexis M Izquierdo (Zaragoza: CIHEAM Centre International de Hautes Etudes Agronomiques Méditerranéennes/HCMR Hellenic National Centre for Marine Research) 63: 158.
25. Taher MM (2007) Effect of fish density and feeding rates on growth and food conversion of gilthead.
26. Yigit M, Bulut M, Ergün S, Güroy D, Karga M, et al. (2012) Utilization of corn gluten meal as a protein source in diets for gilthead sea bream (*Sparus aurata* L.) juveniles. *Fisheries Sciences* 6: 63-73.
27. Lupatsch I, Kissil GWM, Sklan D, Pfeffer E (2001) Effects of varying dietary protein and energy supply on growth, body composition and protein utilization in gilthead sea bream (*Sparus aurata* L.). *Aquaculture Nutrition* 7: 71-80.
28. Bulut M, Yigit M, Ergün S, Kesbic OS, Acar Ü, et al. (2014a) Evaluation of dietary protein and lipid requirements of two-banded seabream (*Diplodus vulgaris*) cultured in a recirculating aquaculture system. *Aquaculture International* 22: 965-973.
29. Rahim A, Abbas G, Ferrando S, Gallus L, Ghaffar A, et al. (2016) Effects of varying dietary protein level on growth, nutrient utilization and body composition of juvenile Black fin seabream, *Acanthopagrus berda* (Forsskal, 1775). *Pakistan Journal of Zoology* 48: 1089-1097.
30. Atienza MT, Chatzifotis S, Divanach P (2004) Macronutrient selection by sharp snout sea bream (*Diplodus puntazzo*). *Aquaculture* 232: 481-491.
31. Tacon AGJ, Cowey CB (1985) Protein and amino acid requirements. In: Tytler P, Calow P, Helm C, editors. *Fish Energetics London & Sydney: New Perspectives*: 155-183.
32. Sabaut JJ, Luquet P (1973) Nutritional requirements of the gilthead bream *Chrysophrys aurata*. Quantitative protein requirements. *Marine Biology* 18: 50-54.
33. Hidalgo F, Alliot E (1988) Influence of water temperature on protein requirement and protein utilization in juvenile sea bass, *Dicentrarchus labrax*. *Aquaculture* 72: 115-129.
34. Kim K, Kayes BT, Amundson HC (1991) Purified diet development and re-evaluation of the dietary protein requirement of fingerling rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 96: 57-67.
35. Espinós FJ, Tomás A, Pérez LM, Balasch J, Jover M (2003) Growth of dentex fingerlings (*Dentex dentex*) fed diets containing different levels of protein and lipid. *Aquaculture* 218: 479-490.
36. Yigit M, Koshio S, Teshima S, Ishikawa M (2004) Dietary protein and energy requirements of juvenile Japanese flounder, *Paralichthys olivaceus*. *J Appl Sci* 4: 486-492.
37. Schuchardt D, Vergara JM, Fernández-Palacios H, Kalinowski CT, Hernández-Cruz CM, et al. (2008) Effects of different dietary protein and lipid levels on growth, feed utilization and body composition of the red porgy (*Pagrus pagrus*) fingerlings. *Aquaculture Nutrition* 14: 1-9.
38. NRC (2011) Nutrient requirements of fish and shrimp. National Research Council. Washington: National Academies.
39. Kaushik SJ, Medale F (1994) Energy requirements, utilization and dietary supply to salmonids. *Aquaculture* 124: 81-97.
40. Vivas M, Rubio VC, Sánchez-Vázquez FJ, Mena C, García BG, et al. (2006) Dietary self-selection in sharpnose seabream (*Diplodus puntazzo*) fed paired macronutrient feeds and challenged with protein dilution. *Aquaculture* 251: 430-437.
41. Ekmann KS, Dalsgaard J, Holm J, Campbell PJ, Skov PV (2013) Effects of dietary energy density and digestible protein:energy ratio on de novo lipid synthesis from dietary protein in gilthead sea bream (*Sparus aurata*) quantified with stable isotopes. *British Journal of Nutrition* 110: 1771-1781.
42. Kesbic OS, Acar Ü, Yigit M, Bulut M, Gültepe N, et al. (2016) Unrefined peanut oil as a lipid source in diets for juveniles of twobanded seabream *Diplodus vulgaris*. *North American Journal of Aquaculture* 78: 64-71.
43. Korkut AY, Balkı D (2004) Effects of different feeding rations on growth of gilthead seabream (*Sparus aurata* L., 1758) in net cages. *Ege Journal of Fisheries and Aquatic Sciences* 21: 235-238.
44. Mongile F, Bonaldo A, Fontanillas R, Mariani L, Badiani A, et al. (2014) Effects of dietary lipid level on growth and feed utilisation of gilthead seabream (*Sparus aurata* L.) reared at Mediterranean summer temperature. *Italian Journal of Animal Science* 13: 30-34.
45. Yigit M, Celikkol B, Bulut M, Decew J, Özalp HB, et al. (2016) Monitoring of trace metals, biochemical composition and growth of Axillary seabream (*Pagellus acarne* Russo, 1827) in offshore copper alloy mesh cages. *Mediterranean Marine Science* 17: 396-403.
46. Bischoff AA, Kube N, Wecker B, Waller U (2005) MARE-Marine artificial recirculated ecosystem: Steps towards closed systems for the production of marine organisms. In: Howell B, Flos R (Eds.). *Lessons from the Past to Optimise the Future*. ESA special publication. 35. European Aquaculture Society. Oostende Belgium 135-136.

47. De La Higuera M, GarcíaGallego M, Sanz A, Hidalgo MC, Suárez MD (1989) Utilization of dietary protein by the eel (*Anguilla anguilla*): optimum dietary protein levels. *Aquaculture* 79: 53-61.
48. Maldonado-García M, Rodríguez-Romero J, Reyes-Becerril M, Álvarez-González CA, Civera-Cerecedo R, et al. (2012) Effect of varying dietary protein levels on growth, feeding efficiency, and proximate composition of yellow snapper *Lutjanus argentiventris* (Peters, 1869). *Latin American Journal of Aquatic Research* 40: 1017-1025.
49. Peres H, Oliva-Teles A (2005) Protein and energy metabolism of European seabass (*Dicentrarchus labrax*) juveniles and estimation of maintenance requirements. *Fisheries Physiology and Biochemistry* 31: 23-31.
50. Green JA, Hardy RW, Brannon EL (2002) The optimum dietary essential:nonessential amino acid ratio for rainbow trout (*Oncorhynchus mykiss*), which maximizes nitrogen retention and minimizes nitrogen excretion. *Fish Physiology and Biochemistry* 27: 109-115.
51. Burel C, Boujard T, Kaushik SJ, Boeuf G, Van Der Geyten S, et al. (2000) Potential of plant-protein sources as fish meal substitutes in diets for turbot (*Psetta maxima*): growth, nutrient utilization and thyroid status. *Aquaculture* 188: 363-382.
52. Fournier V, Huelvan C, Desbruyeres E (2004) Incorporation of a mixture of plant feedstuffs as substitute for fish meal in diets of juvenile turbot (*Psetta maxima*). *Aquaculture* 236: 451-465.
53. Turker A, Yigit M, Ergun S, Karaali B, Erteken A (2005) Potential of poultry by-product meal as a substitute for fishmeal in diets for Black Sea turbot *Scophthalmus maoticus*: Growth and Nutrient Utilization in Winter. *Israeli Journal of Aquaculture Bamidgeh* 57: 49-61.
54. Yigit M, Erdem M, Koshio S, Ergün S, Türker A, et al. (2006) Substituting fish meal with poultry by-product meal in diets for Black Sea turbot *Psettamaeotica*. *Aquaculture Nutrition* 12: 340-347.
55. Harmantepe FB, Yigit M, Dogan G, Karsli Z, Yigit U, et al. (2014) Effects of dietary lipid levels on growth performance and feed utilization in juvenile Black Sea turbot (*Psetta maxima*) with reference to nitrogen excretion. *Marine Science and Technology Bulletin* 3: 21-26.
56. Tulli F, Vachot C, Tibaldi E, Fournier V, Kaushik SJ (2007) Contribution of dietary arginine to nitrogen utilisation and excretion in juvenile sea bass (*Dicentrarchus labrax*) fed diets differing in protein source. *Comparative Biochemistry and Physiology A, MollIntegr Physiol* 147: 179-188.
57. Tulli F, Messina M, Calligaris M, Tibaldi E (2010) Response of European sea bass (*Dicentrarchus labrax*) to graded levels of methionine (total sulfur amino acids) in soya protein-based semi-purified diets. *British Journal of Nutrition* 104: 664-673.
58. Wilson R (1989) Amino acids and proteins. In: J.E. Halver (Ed.) *Fish Nutrition*. San Diego (CA): Academic Press 112-153.
59. Wilson RP (2002) Amino acids and proteins. In: Halver JE & Hardy RW (Ed.). *Fish Nutrition*, 3rd ed. New York: Academic Press 143-179.
60. Stone DAJ (2003) Dietary carbohydrate utilization by fish. *Reviews in Fisheries Science* 11: 337-369.
61. Bibiano JF, Lundstedt LM, Meton I, Baanante IV, Moraes G (2006) Effects of dietary levels of protein on nitrogenous metabolism of *Rhamdia quelen* (Teleostei: Pimelodidae). *Comparative Biochemistry and Physiology A* 145: 181-187.
62. Gaye-Siessegger J, Focken U, Becker K (2006) Effect of dietary protein/carbohydrate ratio on activities of hepatic enzymes involved in the amino acid metabolism of *Nile tilapia*, *Oreochromis niloticus* (L.). *Fish Physiology and Biochemistry* 32: 275-282.
63. Yigit M, Yardim Ö, Koshio S (2002) The protein sparing effects of high lipid levels in diets for rainbow trout (*Oncorhynchus mykiss*, W. 1792) with special reference to reduction of total nitrogen excretion. *Israel Journal of Aquaculture Bamidgeh* 54: 79-88.
64. McClelland G, Zwingelstein G, Weber JM, Brichon G (1995) Lipid composition of tissue and plasma in two Mediterranean fishes, the gilt-head seabream (*Chrysophrys auratus*) and the European seabass (*Dicentrarchus labrax*) *Canadian Journal of Fisheries and Aquatic Sciences* 52: 161-170.
65. Péres H, Gonçalves P, Oliva-Teles A (1999) Glucose tolerance in gilthead seabream (*Sparus aurata*) and European seabass (*Dicentrarchus labrax*). *Aquaculture* 179: 415-423.
66. Pérez-Jiménez A, Hidalgo MC, Morales AE, Arizcun M, Abellan E (2009) Use of different combinations of macronutrients in diets for dentex (*Dentex dentex*) Effects on intermediary metabolism. *Comparative Biochemistry and Physiology A* 152: 314-321.
67. Coutinho F, Peres H, Guerreiro I, Pousão-Ferreira P, Oliva-Teles A (2012) Dietary protein requirement of sharpsnout sea bream (*Diplodus puntazzo*, Cetti1777) juveniles. *Aquaculture* 356: 391-397.
68. Wang F, Han H, Wang Y, Ma X (2013) Growth, feed utilization and body composition of juvenile golden pompano *Trachinotus ovatus* fed at different dietary protein and lipid levels. *Aquaculture Nutrition* 19: 360-367.
69. Li W, Wen X, Zhao J, Li S, Zhu D (2016) Effects of dietary protein levels on growth, feed utilization, body composition and ammonia-nitrogen excretion in juvenile *Nibea diacanthus*. *Fisheries Science* 82: 137-146.
70. Bulut M, Yiğit M, Ergün S, Kesbiç OS, Acar Ü, et al. (2014b) Incorporation of corn gluten meal as a replacement for fish meal in the diets of two banded seabream (*Diplodus vulgaris*) juveniles. *International Journal of Agri Science* 4: 60-65.