

## Case Report

# Mechanisms of Breast Implant Toxicity: Will the Real Ringmaster Please Stand Up

Arthur E Brawer\*

Department of Medicine, Division of Rheumatology, New Jersey, USA

\***Corresponding author:** Arthur E Brawer, Department of Medicine, Division of Rheumatology, 300 Second Avenue, Long Branch, New Jersey 07740, USA. Tel: +17328703133, Fax +17322220824, Email: arthurbrawer@optimum.net

**Citation:** Brawer AE (2017) Mechanisms of Breast Implant Toxicity: Will the Real Ringmaster Please Stand Up. Acad Orthop Res Rheum 1: 104. DOI: 10.29021/2688-9560.100004

**Received Date:** 27 August, 2017; **Accepted Date:** 01 September, 2017; **Published Date:** 08 September, 2017

### Abstract

The past five years have witnessed a recurrent public health debacle encompassing illness caused by silicone gel-filled breast implants. In an attempt to explain this epidemiologic reality, researchers from many institutions have resurrected the faulty device-related autoimmune theories of the early 1990's. These researchers are genuinely convinced that breast implants initiate a novel disease, but their methodology utilized to prove this premise is unsound. The inability to consider mechanisms of widespread biochemical disruptions caused by in vivo exposure to organ silicon molecules is coexisting with multiple deficiencies in detecting the voluminous clinical manifestations of ailing implant recipients. Chief among the latter are the use of truncated data bases to record symptoms and signs, coupled with the lack of appreciation that silicone-induced disease evolves chronologically in a manner simulating a dose-response curve. This failure to correlate the diverse clinical features present in toxic breast implant recipients with logical mechanisms of disease causation provides a clear understanding of why the silicone breast implant chaos of the early 1990's was inevitably destined to repeat itself.

**Keywords:** Biochemistry; Breast implants; Indexing terms; Silicone; Toxicity

### Introduction

Numerous publications in the past five years have supported the assertion that silicone gel-filled breast implants are genuinely capable of causing a novel systemic illness [1-7]. Prior claims of a similar nature existed in the 1990's when controversy first swirled around this issue, but legitimacy at that time was elusive due to faulty methodologies utilized by many investigators [1,4]. Chief among them were the presumptive theories of immune-mediated ailments triggered by these devices. Observations promoting these theories back then included (but were not limited to) (a) the demonstration of extensive inflammatory and immunologic responses in the peri-prosthetic breast tissues and axillary lymph nodes, (b) positive antinuclear antibodies in a third of recipients, (c) the evolution of ever-increasing T cell responsiveness to silica associated with prolonged implantation, (d) the demonstration that silicone gel disseminated to multiple distant sites via the process of gel

bleed through an intact envelope or shell, and (e) case reports of connective tissue diseases related to these devices [8]. Nonetheless, studies in the 1990 have denied the existence of any association between breast implants and autoimmune diseases [9]. Silicone-induced autoimmune theories have recently been resurrected in the form of ASIA (Auto Inflammatory Syndrome induced by adjuvants, or Shoenfeld's syndrome), which promotes the concept of multi-organ inflammatory responses to silicone gel [3,5-7,10]. The deficiencies of ASIA, and the failure of its proponents to consider multiple other mechanisms of disease causation, have been outlined in a recent letter to the editor [11]. In order to further elaborate on these deficiencies one must first understand why the supposedly tightly bound and "Inert" cohesive gel utilized in the new generations of breast implants undergoes dramatic changes in gel properties over time [12]. Breast implants are just one of over 60,000 products composed of organ silicon compounds [13]. Everyday environmental exposure to these myriad products typically winds up in garbage dumps where sludge admixes with dirt and soil. Aerobic and anaerobic bacteria in the soil degrade organ

silicon compounds into molecules of silica (silicon dioxide), silicic acid, silanols, water, and carbon dioxide [14]. It is now known that normal breast tissue is not sterile [15,16]. The diversity of bacteria that are present will routinely colonize the outside and inside of implants once they are inserted into the body [15], whereupon the same degradation processes are initiated [13]. These processes are augmented by hydrolyses released by macrophages, which are also present in abundance outside and inside these devices [8]. In concert with systemic disease onset, the resulting micro dispersion of silanols, silica, silicic acid, and other silicon-containing breakdown products to distant sites and organs occurs long before device rupture [17,18]. Most importantly, this micro dispersion has been documented to exist without concurrent inflammation at these distant sites [19]. Stated another way, in the absence of extensive rupture-induced gel extravasations to the lung or down facial planes, inflammation and immune responses to silicone breast implants are typically present only in the local breast milieu and adjacent axillary nodes (and not at distant sites where excess silicon presence is detected) [20].

## Illness in Implant Recipients

The widespread dispersion of silicon-containing breakdown products causes a myriad of disruptions to the body's biochemistry and enzymatic processes [13,14]. Listed below are examples of how these phenomena correlate with the clinical ailments manifested by implant recipients.

- 75% of my cohort of over 500 implant recipients manifest dry eyes and dry mouth (With markedly abnormal Schirmer tests), but minor salivary (labial) gland biopsies are normal [17]. This phenomenon is likely caused by dysfunction of the proteoglycan receptor for acetylcholine in salivary and lachrymal tissues due to excess deposition of the element silicon in matrix macromolecules [12]. Complaints of dry skin are equally common.
- Although the element silicon forms four bonds like carbon, silicon behaves like a metal at times. Phosphorus in energy systems is metal-ion bound [13]. Therefore, one cannot expect energy production and energy utilization in cells to proceed normally in the presence of excess silicon molecules. Hence, myalgias, fatigue, and weakness occur, but muscle biopsies in my cohort are normal [17].
- Acetylcholine must traverse the matrix at the neuromuscular junction to initiate muscle contraction. Silicon-induced dysfunction of matrix macromolecules can impair this process and also disrupt platelet activation, lymphocyte homing, wound healing, nail growth, and nerve impulse synapse transmission [13].
- The 18,000 square centimeters of skin are quite rich in matrix macromolecules, which explains why splotchy hypo- and hyper-pigmentation, ink blot freckling, multiple nevi, and a variety of skin rashes are common clinical features in ailing implant recipients [17]. In addition, glycosaminoglycans (such as chondroitin sulfate) bind the pre-formed mediators of inflammation in mast cells [13]. Dysfunction of these molecules caused by excess silicon deposition can account for the itching, urticaria, wheezing, and other "Allergic" phenomena experienced by implant recipients. Reactor catalysis is yet another phenomenon that can not only produce skin rashes but also intense burning of the skin [16].
- Synovial fluid analyses from my cohorts' knees reveal less than 1,000 white cells (non-inflammatory). Think of altered matrix macromolecules in cartilage and release of substance P from nerve endings to account for the joint symptoms in implant recipients.
- Silicic acid can cross the blood-brain barrier and chelate catecholamine neurotransmitters (e.g., dopamine) [13]. Hence, cognitive dysfunction occurs in the presence of normal brain MRI scans.
- Silanols can adhere to proteins and enzymes via hydrophobic bonding, with multiple adverse biologic effects [14]. Silanols can also donate a methyl group to mercury (the latter of which may accumulate in one's body from other sources), resulting in the very well-known toxic compound methyl mercury [14].
- Over a dozen heavy metals are utilized in the manufacturing process of silicone gel, and these do not fall out of the soup mixture at the end [20]. These migrate to distant sites along with the silicone breakdown molecules, and can be detected in ailing recipients by appropriate testing. It is therefore no surprise that 40% of my cohorts develop a metallic taste, which resolves after implant removal [21]. Heavy metals adversely affect enzymes, DNA, and leukocyte respiration; can be toxic to muscle, brain, lungs, kidneys, liver, blood, and nerves; and when present in excess amounts can mimic multiple sclerosis, Parkinson's, Alzheimer's, and muscular dystrophy [22]. Clinical correlates reported in implant recipients include (but are not limited to) hair loss, recurrent infections in a variety of sites, weakness, fatigue, muscle twitching, nail cracking and splitting, cognitive dysfunction, a multitude of other nervous system disorders (e.g., parenthesis; dysesthesias; dizziness; gait imbalance); and abdominal pain, nausea, constipation, and loose stools from dysfunction of the myenteric plexus [1-7,13,17,21,23].
- Another cause of recurrent infections in implant recipients (sinus, bladder, pharynx, lung, etc.) is related to immune suppression (not immune stimulation), and occurs from impairment of natural killer cell function [24].
- Ailing silicone recipients also develop odor and smell hypersensitivity. Typical phenomena (which were not present prior

to implantation) include headaches, nausea, and dizziness on exposure to perfumes, room fresheners, cleansers, deodorants, hairsprays, exhaust fumes, and cigarette smoke. Food and pharmaceutical intolerances may also develop. Essentially, when one becomes ill from one toxic entity, small amounts of toxicity coming from elsewhere are poorly tolerated. Most over-the-counter skin products (e.g., moisturizers, skin softeners, wrinkle removers, and cosmetics) also worsen implant-induced phenomena because these products invariably contain silicones and an emulsifier. The latter ingredient enables deep penetration of these products to the subcutaneous fat, where they can become disintegrated (silicones are naturally lipophilic).

- Symptoms such as chest pain and shortness of breath on exertion represent restrictive defects that occur due to local irritation of ribs, intercostal muscles, and intercostal nerves [17]. They often lead to exhaustive cardiopulmonary evaluations which are invariably normal or negative, but nonetheless are erroneously ascribed to serositis of the pleura or pericardium.

## Why Asia is Fantasia (An Improvisation)

In a recent publication Tervaert and colleagues revisit their implant recipients' clinical manifestations and continue to invoke causation via autoimmune and inflammatory mechanisms [6]. This is a gross oversimplification of what is clearly a much more complicated process, and is compounded by a subtle attempt to expand their diagnostic ASIA criteria to be more encompassing (changes that the reader could easily overlook if one did not compare this publication to prior ASIA guidelines) [10]. ASIA goes astray because the diverse inflammatory and immune response to silicone in the local breast environment parallels the development of systemic ailments, but the former does not cause the latter. Stated more simply, the vast majority of silicone-induced ailments emanates from multiple and varied toxic disruptions of the body's biochemistry and have virtually nothing to do with autoimmunity [13,14,22]. Another recent publication by Tervaert and colleagues discusses factors that may contribute to the development of auto antibodies in implant recipients [25]. In actuality, the presence or absence of auto antibodies in patients with these devices has no relevance for diagnosing silicone-induced disease [4,17, 21,23]. In yet another publication Tervaert and colleagues try to prop up their mistakes by publishing a review of comingled conditions (both explicit and/or loosely defined), all of which lack any unifying clinical features but are all supposed to represent silicone-induced toxicity based on a haphazard response to explanation [7]. Their "Kitchen sink" approach is devoid of even the most basic elements of proper investigative methodology including, but not limited to: chronological disease development over time (which has been documented to be a dose-response curve); rupture versus non-rupture analysis; duration of implantation analysis; a more comprehensive, reproducible and consistent data base for recording the ailments of their

patients; prior exchange surgeries; intrinsic validity analysis; explanation results that take into account the duration of prior implantation; and the well-known fact that a patient fulfilling criteria for a classical connective tissue disease doesn't necessarily imply that one has arrived at the correct diagnosis [17,21]. Tervaert and colleagues are genuinely and correctly convinced that silicone gel-filled breast implants are capable of causing a novel systemic illness; however, the methodology of ASIA, and its criteria utilized to diagnose silicone toxicity, is not reliable. The ASIA creators and proponents have reversed the time-honored process inherent to investigations of a new medical disorder. Such a process should initially record detailed observations of patients' ailments at the bedside, followed by subsequent investigations into potential mechanisms of disease causation. Instead, they have placed the cart before the horse by first devising theories of causation, followed by tailoring a set of limited criteria encompassing symptoms and signs that neatly fits their theories. Proponents of ASIA have also abrogated the standard methodology of recording detailed chronological development of multiple clinical phenomena over time, a process (by example) that was crucial in 1975 to the demonstration of Lyme disease as a new illness. The idea that silicone toxicity and other disparate entities encompassed by ASIA are a result of adjuvant-induced immune dysregulation is a concept hatched in a sea of confusion and wishful thinking. One of these other disparate entities deals with illness caused by vaccinations. ASIA was discredited in 2015 by the special masters (judges) in vaccine court in Washington, D.C., and they will no longer allow it to be referenced by any experts testifying on behalf of vaccine injury claimants. Still another one of these disparate entities deals with Gulf War syndrome in 1991. Seven years later, in 1998 in a publication in JAMA, this disorder was proven to be an epidemiological reality, but it had virtually nothing to do with autoimmunity [26]. If ASIA becomes the standard by which silicone toxicity is defined, it is quite likely that future scientific panels will negate this as well. This, in turn, may well bury the validity of breast implant illness a second time, forever dooming the legions of ailing women from achieving any vindication.

## Case Studies

The following two case histories exemplify the recognition and vindication sought by ailing implant recipients.

Case 1: A 22 y.o.w.f., previously in good health and on no medications, underwent bilateral breast augmentation in 2013 with the insertion of Mentor cohesive silicone gel-filled breast implants. Within two months she began to experience breast pains, chest pain and pressure, fatigue, weakness, and bilaterally enlarged axillary lymph nodes. Over the next three years she developed a multisystem illness encompassing night sweats, metallic taste, headaches, vitiligo on her arms, skin freckling on her arms and anterior chest wall, foul body odor, cognitive dysfunction (memory lapses, poor assimilation of new material, confusion, and problems with

name recall and word recall), pain and stiffness in her hands and knees, two hours of morning stiffness, photosensitivity, nail cracking and splitting, two different types of skin rashes on the anterior chest, posterior trunk, shoulders, arms and feet (Punctate red pin-head sized papules; also pustules), itching, poor wound healing, bruising, shortness of breath on exertion, dry eyes, abdominal pain, loose stools alternating with constipation, pins and needles and burning pains in her extremities, sore throats, dysphasia, and odor and smell hypersensitivity (with nausea, dizziness, and headaches from cleansers, make-up, perfumes, detergents, room fresheners, deodorants, cigarette smoke, and exhaust fumes). Physical examination was notable for 0mm tear formation on a Schirmer test, 1.5cm bilateral axillary lymph nodes, freckling and pigment changes on the skin (as noted above), tenderness to palpation of multiple anterior ribs, swelling with pain on motion in the PIP joints of her hands, micronodularity of both breasts, two types of skin rashes (as noted above), nail friability, and easy muscle fatigue on repetitive muscle testing. Routine laboratory tests, as well as ANA, were normal or negative. Removal of intact implants and capsulectomies were performed 3 and ½ years after her original surgery. Within seven weeks she began to notice some improvement in her condition, and six months after explanation she has experienced improvement and/or resolution of 80% of her overall symptoms and signs.

**Case 2:** A 40 y.o.w.f., previously in good health and on no medications, underwent bilateral breast augmentation in 2012 with the insertion of Mentor cohesive silicone gel-filled breast implants. Within one month she began to experience breast pains, itching, urticaria on her abdominal wall, and red pin-head sized papules on her neck and arms. Over the next four years she developed a multisystem illness encompassing dry eyes and dry mouth, chest pain, bilateral axillary lymphadenopathy, sore throats, recurrent bladder infections, recurrent ear infections, fatigue, muscle weakness, pain and stiffness in her shoulders, hands, wrists, knees, hips, and ankles, one hour morning stiffness, metallic taste, myalgias, abdominal pain, nausea and vomiting, loose stools, night sweats, muscle twitching, cognitive dysfunction (Similar to case 1), pins and needles and burning pains in her extremities, blurry vision, photosensitivity, nail cracking and splitting, headaches, bruising, shortness of breath on exertion, splotchy hyper pigmentation of her face and neck, hypo pigmentation on her anterior chest, posterior trunk, shoulders and arms, foul body odor, dysphasia, and odor and smell hypersensitivity (similar to case 1). Physical examination was notable for 0mm tear formation on a Schirmer test, 2cm bilateral axillary lymph nodes, pigment changes on the skin (as noted above), two types of skin rashes (as noted above), tenderness to palpation of multiple anterior ribs, pain on motion of multiple small and large joints, and several ecchymoses. Routine laboratory tests, as well as an ANA were normal or negative. A rheumatologist prescribed NSAID's, Methotrexate, Leflunamide, and Enbrel without help. Removal of intact implants and capsulectomies were

performed 4 and ½ years after her original surgery. Within four weeks she began to notice some improvement in her condition, and eight months after explanation she has experienced improvement and/or resolution of 90% of her overall symptoms and signs.

## Other Observations

Recent publications by University of Oregon researchers have added still another dimension to this saga [27,28]. It is alleged that silicone gel can act as a magnet for certain chemicals and toxins that one might encounter in their everyday environment. It is already known that silicones attract estrogen molecules, thereby acting as an independent reservoir to disturb the body's equilibrium of normally well-controlled hormone levels [13]. Silicones are also lipophilic, and they readily adhere to fatty acids and phospholipids (Including myelin components) as well as adipocytes [27,28]. The chemicals and toxins studied by the Oregon researchers are also lipophilic. Translation: chemicals and toxins are therefore naturally attracted to silicones and vice versa. In addition to heavy metals, numerous chemicals are utilized in the manufacturing processes to synthesize silicone gel [20]. Many of these are neurotoxins and carcinogens, and given the recent revelations above they probably do not fall out of the soup mixture at the end. In addition to providing still another mechanism for silicone toxicity, could these phenomena also be related to the recent evidence linking a rare type of cancer, Anaplastic Large Cell Lymphoma (ALCL), to certain types of silicone gel-filled breast implants [29]? Why doesn't every recipient of silicone gel-filled breast implants become ill? Part of the answer may lie in the human evolutionary process itself. An enzyme, silicase, is normally found in invertebrates such as starfish species, and aids in the degradation and dissolution of silica and other harmful silicon-containing molecules [30]. This enzyme activity was lost in higher organisms [13], but perhaps there are some women who still retain the capacity to synthesize it, and thus have the capability to neutralize silicon-containing compounds. If there is a protective lineage, then such recipients would be considered "Non-Responders," as opposed to the "Over-Responders" hypothesized by the ASIA proponents who opine that one's genetic make-up is a factor producing adverse systemic autoimmune reactions to silicone [5].

## Conclusions

So, what should we call this new disease entity? ASIA? Human adjuvant disease? Silicone implant incompatibility syndrome? Autoimmune Systemic Silicone intolerance? Siliconosis? For simplicity, we probably should use the phrase "Silicone-induced Toxicity." But clearly there is nothing "Simplistic" regarding this entity, and scientists with a PhD in biochemistry need to get involved with the research. The absence of their participation in the early 1990's, in part because of the tunnel vision of rheumatologists and immunologists, significantly contributed to the inevitable recurrence of today's silicone breast implant public health debacle

[1,4,13]. It is now perfectly clear that silicone gel-filled breast implants cause a distinct and verifiable illness if one utilizes proper methodology to define it [1,4,17,21].

## References

1. Brawer A (2012) Bones, groans, and silicone. *Lupus* 21: 1155-1157.
2. Majers MC, deBlok CJM, Niessen FB, van der Veldt AA, Ritt MJ, et al. (2013) Women with silicone breast implants and un-explained systemic symptoms: A descriptive cohort study. *Netherlands J Med* 71: 534-540.
3. Colaris MJL, deBoer M, van der Hulst RR, Tervaert JWC (2017) Two hundred cases of ASIA Following silicone implants: a comparative study of 30 years and a review of current Literature. *Immunol Res* 65: 120-128.
4. Brawer A (2017) Destiny rides again: the reappearance of silicone gel-filled breast implant toxicity. *Lupus* 26: 1060-1063.
5. Watah A, Quaresma M, Brown S, Tervaert JW, Rodríguez-Pint I5, et al. (2017) Autoimmune/inflammatory syndrome induced by adjuvants (Shoenfeld's syndrome)-An update. *Lupus* 26: 675-681.
6. Tervaert JWC, Colaris MJ, van der Hulst RR (2017) Silicone breast implants and autoimmune Rheumatic disease: myth or reality. *Curr Opin Rheumatol* 29: 348-354.
7. deBoer M, Colaris M, van der Hulst RRWJ, Tervaert JWC (2017) Is explanation of silicone breast Implants useful in patients with complaints? *Immunol Res* 65: 25-36.
8. Shanklin DR, Smalley DL (1998) the immunopathology of siliconosis. *Immunol Res* 18: 125-173.
9. Gabriel SE, O'Fallon WM, Kurland LT, Beard CM, Woods JE, et al. (1994) Risk of connective tissue diseases and other disorders after breast implantation. *N Engl J Med* 330: 1697-1702.
10. Cervera R (2011) ASIA: A new systemic autoimmune syndrome? *Lupus* 20: 665-666.
11. Brawer A (2017) Autoinflammatory Syndrome Induced by Adjuvant (ASIA) syndrome is misguided. *Natural Med J* 9: 5.
12. Bodin F, Jung C, Dieval F, Chakfe N, Wisniewski S, et al. (2015) Aging of retrieved gel breast implants: A comparison between two product generations. *J MechanBehav Biomed Materials* 46: 11-22.
13. Brawer A (1998) Silicon and matrix macromolecules: New research opportunities for old diseases from analysis of potential mechanisms of breast implant toxicity. *Med Hypoth* 51: 27-35.
14. Hirner AV, Flassbeck D, Gruemping R (2003) Organosilicon Compounds in the Environment, in Peter J. Craig, editor, *Organometallic Compounds in the Environment*, 2<sup>nd</sup> edition Chapter 8: 305-351.
15. Ajdic D, Zoghbi Y, Gerth MS, Panthaki ZJ, Thaller S (2016) The relationship of bacterial biofilms and capsular contracture in breast implants. *Aesthetic Surg J* 36: 297-309.
16. Hiekin TJ, Chen J, Hoskin TL, Walther-Antonio M, Johnson S, et al. (2016) The microbiome of aseptically collected human breast tissue in benign and malignant disease. *Scientific Reports* 3: 30751.
17. Brawer A (1996) Chronology of systemic disease development in 300 symptomatic recipients of Silicone gel-filled breast implants. *J Clean Technol, Environ Toxicol and Occup Med* 5: 223-233.
18. Brawer A (1996) Clinical features of local breast phenomena in 300 symptomatic recipients of silicone gel-filled breast implants. *J Clean Technol, Environ Toxicol and Occup Med* 5:235-247.
19. Kappel RM, Boer LL, Dijkman H (2016) Gel bleed and rupture of silicone breast implants investigated by light-, electron microscopy and energy dispersive x-ray analysis of internal organs and nervous tissue. *Clin Med Reviews and Case Reports* 3: 2378-3656.
20. Food and Drug Administration Summary of Safety and Effectiveness Data November 17 2006 (PMA P030053).
21. Brawer A (2000) Amelioration of systemic disease following removal of silicone gel-filled breast implants. *J Nutrit& Environ Med* 10: 125-132.
22. Jaishankar M, Tseten T, Krishnamurthy N (2014) Toxicity, mechanism, and health effects of some heavy metals. *Interdiscip Toxicology* 7: 60-72.
23. Borenstein D (1994) Siliconosis: A spectrum of illness. *Semin Arthritis Rheum* 24: 1-7.
24. Campbell A, Brautbar N, Vojdani A (1994) Suppressed natural killer cell activity in patients with silicone breast implants: reversal upon explantation. *Toxicol Ind Health* 10: 149-154.
25. Colaris MJL, vanderHulst, RR, Tervaert JWC (2017) Vitamin D deficiency as a risk factor for the development of autoantibodies in patients with ASIA and silicone breast implants: a cohort study and review of the literature. *Clinical Rheumatol* 36: 981-993.
26. Fukuda K, Nisenbaum R, Stewart G, Robin L, Washko RM, et al. (1998) Chronic multisystem illness affecting air force veterans of the Gulf War. *JAMA* 280: 981-988.
27. O'Connell SG, Kincl LD, Anderson KA (2014) Silicone wristbands as personal passive samplers. *Environ Science & Technol* 48: 3327-3335.
28. O'Connell SG, Kerkvliet NI, Carozza S, Rohlman D, Pennington J (2015) et al. In vivo partitioning to silicone implants: Implications for use in biomonitoring and body burden. *Environ International* 85: 182-188.
29. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, et al. (2016) The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood* 127:2375-2390.
30. Schroder HC, Krasko A, LePennec G, Adell T, Wiens M, et al. (2003) Silicase, an enzyme which degrades bigamous amorphous silica: contribution to the metabolism of silica deposition in the demosponge *Subefritesdomuncula*. *ProgMol Sub cell Biol* 33: 249-268.