

# Phylogenetic Analysis of Metalloprotease from Transcriptome of Venom Gland of *Hemiscorpius lepturus*

Fatemeh Kazemi-Lomedashta\*, Delavar Shahbazzadeha, Mahdi Behdania

Venom & Biotherapeutics Molecules Laboratory, Biotechnology Research Center, Pasteur Institute of Iran, Iran

**\*Corresponding author:** Fatemeh Kazemi-Lomedasht, Venom & Biotherapeutics Molecules Laboratory, Biotechnology Research Center, Pasteur Institute of Iran, Iran. Email: fa\_kazemi@pasteur.ac.ir

**Citation:** Kazemi-Lomedashta F, Shahbazzadeha D, Behdania M (2018) Phylogenetic Analysis of Metalloprotease from Transcriptome of Venom Gland of *Hemiscorpius lepturus*. J Trop Med Health: JTMH-117. DOI: 10.29011/JTMH-117. 000017

**Received Date:** 27 January, 2018; **Accepted Date:** 23 April, 2018; **Published Date:** 30 April, 2018

## Abstract

*Hemiscorpius lepturus* is a dangerous scorpion and referred to health concern issue in Khuzestan, Iran. The venom of *H. lepturus* is cytotoxic and its effect is similar to spider *Loxosceles reclusa*. Metalloproteinases are the important class of enzymes in the venom that has hemorrhagic activity. The early finding suggests the existence of metalloproteases in the transcriptome of venom gland of *H. lepturus*. Phylogenetic analysis was accomplished to reveal the evolutionary relationship of identified metalloproteases. The phylogenetic tree was constructed by Molecular Evolutionary Genetics Analysis software and neighbor-joining method. Results showed among three sequences, two metalloproteinases named HLMP1 and HLMP3 of *H. lepturus* were most close to spider *P. tepidarioru*. The third sequence named HLMP2 was different and formed an independent clade in the phylogenetic tree. The results suggest that the sequence of metalloproteases in the venom component of *H. lepturus* is similar to the spider than the scorpion.

**Keywords:** *Hemiscorpius lepturus*; Iranian Scorpion; Metalloproteinases; Phylogeny; Venom Component

## Introduction

*Hemiscorpius lepturus* is one of the dangerous scorpions of Iran. The pathological effects of *H. lepturus* are different of other scorpions and results in failure of kidney, liver as well as blood cells [1]. Various studies have evaluated the venom component of *H. lepturus* and various toxic peptides and proteins of *H. lepturus* have been identified like: hemitoxin [2], hemicalcin [3], heminecrolysin [4] and hemilipin [5]. Transcriptome analysis of venom gland of *H. lepturus* was performed in previous study and whole venom component of this scorpion was identified [6]. Early finding of transcriptome analysis suggests existence of metalloprotease enzyme in the venom gland of *H. lepturus* [6,7]. However, the metalloprotease activity of *H. lepturus* has been evaluated using zymography analysis [8]. Matrix metalloproteinases (MMPs) play a role in growth and cell differentiation through direct and indirect mechanism on growth factors function [9,10,11]. It seems that MMP can affect cellular attachment to the matrix by proteolysis of the adhesion sites [12]. In the early stage, MMP plays a role

in cell proliferation and cell survival of tumor cells [13]. There is diversity in function of various metalloproteinases of different organisms [14]. Metalloproteases are important enzymes in the venom component. They play important role in hemorrhagic activity of venom [15]. A metalloproteinase named, BumaMPs1, has been identified in the venom of *Buthus martensi* scorpion [15]. BumaMPs1 characterized as independent metalloproteinase of others [15].

Phylogenetic analysis results showed that BumaMP1 is most similar to *Mesobuthus eupeus* scorpion metalloproteinase [15]. Here, phylogenetic analysis of the metalloproteinase (identified in transcriptome of venom gland) of *H. lepturus* was performed for the first time to identify evolutionary relationship of the enzyme in three different taxa: Insecta, Ophidia and Arachnida.

## Materials and Methods

### Metalloproteinase Sequences

The transcriptome analysis of venom gland of *Hemiscorpius lepturus* was accomplished in previous work [6]. Metalloproteinase sequences identified in the transcriptome and submitted to NCBI

as follows accession numbers: HLMP1; KX924496, HLMP2; KX924497, HLMP3; KX924498 [6,7]. The protein sequences were aligned using Basic Local Alignment Search Tool (BLAST) of National Center for Biotechnology Information (NCBI). The BLASTp tool was used and alignment was performed in Non-redundant protein sequences (nr databank) (<http://nlm.nih.gov/BLAST/>).

### Phylogenetic Analysis

To construct phylogenetic tree, the sequences with high similarity to the metalloprotease sequences of *H. lepturus* were selected from three taxa: Arachnida (*Latrodectus hesperus* ADV40108; *Tityus serrulatus* CDJ26716.1; *Stegodyphus mimosarum* KFM63177.1; *Parasteatoda tepidariorum* XP\_015909675.1, XP\_015919212.1, XP\_021003544.1 ; *Tityus serrulatus* CDJ26717.1 ; *Ixodes scapularis* XP\_002406806.1 , XP\_002435824.1), Ophidia (*Thamnophis sirtalis* XP\_013907739.1 ; *Protobothrops mucrosquamatus* XP\_015680512.1 ; *Python bivittatus* XP\_007429967.1) and Insecta (*Pogonomyrmex barbatus* XP\_011632701.1 ; *Pediculus humanus corporis* XP\_002426660.1 ; *Zootermopsis nevadensis* XP\_021917577.1 ; *Stomoxys calcitrans* XP\_013099168.1 ; *Wasmannia auropunctata* XP\_011692954.1, XP\_011704691.1 ; *Halyomorpha halys* XP\_014273620.1 ; *Aedes albopictus* XP\_019532310.1 ; *Camponotus floridanus* XP\_011253904.1 ; *Helicoverpa armigera* XP\_021196322.1 ; *Rhagoletis zephyria* XP\_017463716.1 ). The polygenetic tree was constructed by Molecular Evolutionary Genetics Analysis software (MEGA5) using neighbor-joining method and 1000 bootstrap replicate [16,17].

## Results


### Alignment Analysis

Alignment of HLMP1 in nr databank of NCBI showed 50, 51, 53 and 56 percent identity with astacin-like metalloprotease, [*Latrodectus hesperus* ADV40108.1], astacin-like metallopeptidase 1 protein [*Tityus serrulatus* CDJ26716.1], Zinc metalloproteinase nas-4, partial [*Stegodyphus mimosarum* KFM63177.1], astacin-like metalloprotease toxin5[*Parasteatoda tepidariorum* XP\_015909675.1, XP\_015909677.1], respectively. Alignment results of HLMP2 showed 35, 36 and 45 percent identity to metalloprotease1 [*Ixodes persulcatus* AIE44747.1], metalloprotease [*Ixodes scapularis* XP\_002435641.1], metalloserrulase 19 [*Tityus serrulatus* AM002561.1], respectively. Alignment analysis of HLMP3 showed 50, 51, 53 and 56 percent identity to astacin-like metalloprotease [*Latrodectus hesperus* ADV40108.1], astacin-like metallopeptidase 1 protein [*Tityus serrulatus* CDJ26716.1], Zinc metalloproteinase nas-4 [*Stegodyphus mimosarum* KFM63177.1], astacin-like metalloprotease toxin 5 [*Parasteatoda tepidariorum* XP\_015909675.1, XP\_015909677.1], respectively.

### Phylogenetic Analysis

Phylogenetic tree was constructed using 23 sequences of three different taxa: Insecta (11 sequences), Ophidia (3 sequences) and Arachnida (9 sequences). There were different species in Arachnida taxa: scorpions (2 sequences), spiders (5 sequences) and tick (2 sequences). As it can be seen in (figure 1) HLMP1 and HLMP3 organized a single clade close to *Parasteatoda tepidariorum* (XP\_021003544.1) and HLMP2 was in other clade.

The results indicate that metalloprotease sequences of *H. lepturus* scorpion are similar to *P. tepidariorum* spider.



**Figure 1:** Phylogenetic tree. The phylogenetic tree was constructed by MEGA 5 using neighbor-joining method and 1000 bootstrap replicate. A: Orginal tree, B: Bootstrap consensus tree. HLMP1 and HLMP3 organized a single calde) close to *Parasteatoda tepidariorum* (XP\_021003544.1). HLMP2 formed an independent clade.

## Discussion

Metalloproteinases exist in the venom of different venomous animal like snake and scorpion [8,15,18,]. Hemorrhagic activity of venom is related to the presence of enzymes like metalloproteinase in the venom [15]. The metalloproteinases fall into different groups depending on their metal atom [19]. Three sequences in the transcriptome of *H. lepturus* are identified that showed similarity to metalloproteinase sequences. HLMP1 and HLMP3 showed similarity with Astacin-like metalloproteinase toxin but HLMP2 showed similarity with venom metalloproteinase antarease. Antarease-like Zinc-metalloproteinases in the venom of scorpion *Tityus serrulatus* leads to acute pancreatitis [20]. Targeting of metalloproteinase in the scorpion venom is novel anti-venom strategy [20]. In phylogenetic analysis of three metalloproteinases from transcriptome of venom gland of *H. lepturus*, two clade was organized. The HLMP1 and HLMP3 were in one clade and HLMP2 was in independent clade. The results indicating difference of these sequences with each other. The analysis was performed on the sequences that achieved by transcriptome analysis of venom gland. It has been reported that some sequences in the transcriptome may not detect in the proteome and some sequences in the proteome may not predict in the transcriptome [21]. Therefore, it seems necessary to compare transcriptome and proteome of *H. lepturus* venom at same time to analyze the accuracy of achieved sequences.

## Conclusion

Here, for the first-time phylogenetic analysis of three metalloproteinase sequences from transcriptome of venom gland of *H. lepturus* was performed. Results showed that two metalloproteinases (HLMP1 and HLMP3) of *H. lepturus* are most close to spider *P. tepidariorum*. However, the sequence of HLMP2 was differ from HLMP1 and HLMP3 and formed an independent clade in phylogenetic tree. However further studies on proteome of *H. lepturus* is needed.

## Acknowledgment

This study was supported by Pasteur Institute of Iran, Tehran, Iran. The authors warmly thanks from Pasteur Institute of Iran, Tehran, Iran for supporting current study.

## Conflict of interest

There is no conflict of interest.

## References

1. Jalali A, Pipelzadeh MH, Sayedian R, Rowan E (2010) A review of epidemiological, clinical and *in vitro* physiological studies of envenomation by the scorpion *Hemiscorpius lepturus* (Hemiscorpiidae) in Iran. Toxicon 55: 173-179.
2. Srairi-Abid N, Shahbazzadeh D, Chatti I, Mlayah-Bellalouna S, Mejdoub H, et al. (2008) Hemitoxin, the first potassium channel toxin from the venom of the Iranian scorpion *Hemiscorpius lepturus*. The FEBS journal 275: 4641-4650.
3. Shahbazzadeh D, Srairi-Abid N, Feng W, Ram N, Borchani L, et al. (2007) Hemicalcin, a new toxin from the Iranian scorpion *Hemiscorpius lepturus* which is active on ryanodine-sensitive  $Ca^{2+}$  channels. Biochemical journal 404: 89-96.
4. Borchani L, Sassi A, Shahbazzadeh D, Strub JM, Tounsi-Guettet H, et al. (2011) Heminecrosis, the first hemolytic dermonecrotic toxin purified from scorpion venom. Toxicon 58: 130-139.
5. Jridi I, Catacchio I, Majdoub H, Shahbazzadeh D, El Ayeb M, et al. (2015) Hemilipin, a novel *Hemiscorpius lepturus* venom heterodimeric phospholipase A2, which inhibits angiogenesis *in vitro* and *in vivo*. Toxicon 105: 34-44.
6. Kazemi-Lomedasht F, Khalaj V, Bagheri KP, Behdani M, Shahbazzadeh D (2017) The first report on transcriptome analysis of the venom gland of Iranian scorpion, *Hemiscorpius lepturus*. Toxicon 125: 123-130.
7. Jahdasani R, Jamnani FR, Behdani M, Habibi-Anbouhi M, Yardehnavi N, et al. (2016) Identification of the immunogenic epitopes of the whole venom component of the *Hemiscorpius lepturus* scorpion using the phage display peptide library. Toxicon 24: 83-93.
8. Seyedian R, Pipelzadeh MH, Jalali A, Kim E, Lee H, et al. (2010) Enzymatic analysis of *Hemiscorpius lepturus* scorpion venom using zymography and venom-specific antivenin. Toxicon 56: 521-525.
9. Kazushi I, Hiramatsu A, Fukushima D, Pierschbacher MD, Okada Y (1997) Degradation of decorin by matrix metalloproteinases: identification of the cleavage sites, kinetic analyses and transforming growth factor- $\beta$ 1 release. Biochemical Journal 322: 809-814.
10. Buckley A (2016) Potential Therapeutic Efficacy of a Novel Metalloproteinase Inhibitor, Extracellular Matrix Protection Factor 1, in Human Osteoarthritic Chondrocyte Primary Cultures.
11. Kim BJ, Hur JW, Park JS, Kim JH, Kwon TH, et al. (2016) Expression of matrix metalloproteinase- 2 and- 9 in human ligamentum flavum cells treated with tumor necrosis factor-  $\alpha$  and interleukin-1 $\beta$ . Journal of Neurosurgery: Spine 24: 428-435.
12. Giannelli G, Falk-Marzillier J, Schiraldi O, Stettler-Stevenson WG, Quaranta V (1997) Induction of cell migration by matrix metalloproteinase-2 cleavage of laminin-5. Science 277: 225-228.
13. Rudolph-Owen LA, Chan R, Muller WJ, Matrisian LM (1998) The matrix metalloproteinase matrilysin influences early-stage mammary tumorigenesis. Cancer Research 58: 5500-5506.
14. Azofeifa-Cordero G, Arce-Estrada V, Flores-Díaz M, Alape-Girón A (2008) Immunization with cDNA of a novel P-III type metalloproteinase from the rattlesnake *Crotalus durissus durissus* elicits antibodies which neutralize 69% of the hemorrhage induced by the whole venom. Toxicon 52: 302-308.
15. Xia X, Ma Y, Xue S, Wang A, Tao J, et al. (2013) Cloning and molecular characterization of BumaMPs1, a novel metalloproteinases from the venom of scorpion *Buthus martensi* Karsch. Toxicon 76: 234-238.
16. Zuckerkandl E, Pauling L (1965) Evolutionary divergence and convergence in proteins. Evolving genes and proteins 97: 97-166.

17. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, et al. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular biology and evolution* 28: 2731-2739.
18. Fox JW, Serrano SM (2008) Insights into and speculations about snake venom metalloproteinase (SVMP) synthesis, folding and disulfide bond formation and their contribution to venom complexity. *The FEBS journal* 275: 3016-3030.
19. Hooper NM (1994) Families of zinc metalloproteases. *FEBS letters* 354: 1-6.
20. Ortiz E, Rendón-Anaya M, Rego SC, Schwartz EF, Possani LD (2014) Antarease-like Zn-metalloproteases are ubiquitous in the venom of different scorpion genera. *Biochimica et Biophysica Acta (BBA)-General Subjects* 1840: 1738-1746.
21. Shan LL, Gao JF, Zhang YX, Shen SS, He Y, et al. (2016) Proteomic characterization and comparison of venoms from two elapid snakes (*Bungarus multicinctus* and *Naja atra*) from China. *Journal of proteomics*, 138: 83-94.