



Research Article

Resection Margins in Squamous Cell Lip Carcinoma According to the Tumor Site and Size

Cristina Stanescu*, Georgiana-Daniela Stanescu, Camelia Tamas

Faculty of Medicine and Pharmacy "Dunărea de Jos" University of Galati, Romania

University of Medicine and Pharmacy "Gr. T. Popa" Iasi, Romania

Emergency County Hospital of Braila, Romania

***Corresponding author:** Stanescu C. Faculty of Medicine and Pharmacy "Dunărea de Jos" University of Galati, University of Medicine and Pharmacy "Gr. T. Popa" Iasi, Emergency County Hospital of Braila, Romania.

Citation: Stanescu C, Stanescu GD, Tamas C (2023) Resection Margins in Squamous Cell Lip Carcinoma According to the Tumor Site and Size: Research Article. *Plast Surg Mod Tech* 7: 166. DOI: 10.29011/2577-1701.1000166

Received Date: 01 May, 2023; **Accepted Date:** 09 May, 2023; **Published Date:** 12 May, 2023

Abstract

Introduction: Lip cancer is the most frequent malignant tumors of the oral cavity. Lip carcinoma could appear on pre-cancerous lesions (such as radiodermatitis, chronic cheilitis, xeroderma pigmentosum etc.) and is related with other risk factors like smoking, chronic alcohol consumption, immunosuppression, HPV infection etc. **Materials and Methods:** This study was conducted on a representative sample made of 164 patients diagnosed with squamous cell carcinoma and 62 patients with squamous cell lip carcinoma from 2010 to 2018 at the Plastic Surgery Compartment at the Emergency County Hospital of Braila. The incidence of squamous cell lip carcinoma increase in the last years from 20% in 2010 to 26% in 2018. This study analyses clinical characteristics and histopathological types of lip carcinoma, frequency of the squamous cell lip carcinoma, the relation between the resection margins and tumor size, types of local flaps used in the reconstruction of consecutive defects, local evolution and possible metastatic spread to the lymph node of the neck. Squamous cell lip carcinoma was located to the lower lip at 40 patients, to the commissure at 14 patients and to the upper lip at 8 patients. The youngest patient with squamous cell lip carcinoma was 22 years old. **Results and Discussion:** Lesions smaller than 1 cm were more frequent (62%) compare to the lesions which involved more than half of the lip (8%). For small tumors, under 1 cm, the resection margins was made at 4 mm, for tumors with sizes between 1-2 cm the tumor resection was performed 6 mm from the tumor margins and for large tumors, more than 2 cm, the resection margins increase to 1 cm. Early stage tumors were treated by direct suture with a very good prognosis. For advanced lesions we used a variety of flaps for lip reconstruction: Gillies flap, Abbé flap, Karapandzic flap, Mc Gregor flap, Estlander flap, nasolabial flaps (Fujimori and Ombredane) etc. Two patients return for commissuroplasty. We used a Converse flap and a Zisser flap. **Conclusions:** The successful reconstruction depends on preoperative planning according to the tumor site and size. Lymph node metastases significantly reduce long term survival. For a plastic surgeon the aesthetic appearance has the same importance like the functional results.

Keywords: Squamous Cell Lip Carcinoma; Incidence; Risk Factors; Resection Margins; Surgical Reconstruction; Survival Rate.

Introduction

Lip cancer is the most frequent malignant tumors of the oral cavity [1]. The squamous cell carcinoma is the most frequent tumor in the lower lip, on the red part, whereas the basal cell carcinoma usually involves the white upper lip. The lesion caused

by squamous cell lip carcinoma is often asymptomatic and the clinical appearance is highly variable from a reddish skin plaque to an ulcerated lesion with hard raised edges. Squamous cell lip carcinoma are slow-growing, but occasionally, these cancers can have a behaviour more aggressively, spreading to regional lymph nodes at a relatively early stage [2]. The incidence in rural patients is higher compare to urban patients because of their outdoors activities. Long term exposure to sun radiation is directly involved in the pathogeny of facial cutaneous carcinoma [3]. Lip carcinoma

could appear on pre-cancerous lesions, such as radiodermatitis, chronic cheilitis, xeroderma pigmentosum etc. Most skin cancers are not caused by an inherited faulty gene that can be passed on to the next generation, but members of families have the same skin type, which may increase their risk of developing a skin cancer. Lip carcinoma is related with other risk factors like smoking, chronic alcohol consumption, immunosuppression, HPV infection etc [4]. Another possible risk factors are represented by over exposure to some chemicals such as coal tar, soot, pitch, asphalt, petroleum derivatives, paraffin waxes, arsenic etc [5].

Excision is undertaken with 4 mm margins for low risk and more than 6 mm for high risk lesions [6]. There is no randomised controlled study to show that Mohs technique is superior to conventional surgery [7]. Regarding the choice for lip reconstruction after wide excision, several issues should be considered including contour, texture, colour, aesthetic units, donor site and the wishes of the patients. Early stage tumors treated by direct suture have a very good prognosis. For advanced lesions a variety of flaps could be used, according to the preoperative plan [8]. The percentage of patients with lymph node metastases increases considerably in advanced stage tumor. Lymph node metastases significantly reduce long term survival [9].

Materials and Methods

This study was conducted on a representative sample made of 164 patients diagnosed with squamous cell carcinoma and 62 patients with squamous cell lip carcinoma from 2010 to 2018 in the Plastic Surgery Compartment at the Emergency County Hospital of Braila. This study analyses clinical characteristics and histopathological types of lip carcinoma, frequency, the relation between the resection margins and tumor size, types of local flaps used in the reconstruction of consecutive defects, functional results, local evolution and possible metastatic spread to the lymph node of the neck. We quantified the characteristic parameters represented by age, sex, residence, types of carcinoma, anatomical areas. The incidence in rural patients is high compare to the urban patients. The incidence in male patients is higher (58%) than female (42%). Squamous cell lip carcinoma was located to the lower lip at 40 patients (64.5%), to the commissure at 14 patients (22.5%) and to the upper lip at 8 patients (13%). The youngest patient with squamous cell lip carcinoma was 22 years old. Eight patients developed two types of cutaneous carcinoma at the same time in different areas.

Lesions smaller than 1 cm were more frequent (62%), compare to the lesions which involved more than half of the lip

(8%). 96% of squamous cell lip carcinoma involve the vermillion. The successful of reconstruction depends on preoperative planning according to the tumor site and size. For small tumors, under 1 cm, the resection margins was made at 4 mm, for tumors with diameter between 1-2 cm the tumor resection is performed 6 mm from the tumor margins and for large tumors, more than 2 cm, the resection margins increase to 1 cm. Numerous techniques have been developed for lip reconstruction. The choice depend on the position of tumor and its extension level. Early stage tumors was treated by direct suture after V-shaped excision, with a very good prognoses. For advanced lesions we used a variety of flaps for lip reconstruction: Gillies flap, Mc Gregor flap, Karapandzic flap, Abbé flap, Estlander flap, nasolabial flap (Fujimori and Ombredane) etc. Two patients return for commissuroplasty. We used a Converse flap and a Zisser flap. Suture must consider all anatomical planes, using separate absorbable threads for mucosa and muscular plane and nonabsorbable threads for skin. Lymph node metastases significantly reduce long term survival.

Results and Discussion

The distribution of patients with squamous cell carcinoma has revealed that the highest incidence of this type of cancer is recorded in person aged over 50 years. The incidence of squamous cell lip carcinoma increase in the last years from 20% in 2010 to 26% in 2018. Local flaps was the first option for the most patients with large tumors which requiring postoperative reconstructions. The reconstruction of postoperative defects depend on the position of the lesion and its extension level. The patients with early stage tumors recovered within 3 weeks after surgery with good long term aesthetic and functional results. Smaller lesions with diameter under 1 cm were most frequent (62%). In a few cases (8%) the lesions involved more than two-thirds of the lip. In 78% of cases a direct suture was performed, after V-shaped excision (Figure 1, 2).



Figure 1: 74 Year Old Woman with Lower Lip Carcinoma



Figure 2: Direct Suture After V-shaped Excision

In 14% of cases the lip reconstruction required the use of a single flap (Figure 3, 4). For near total lip defects (8% of cases) it was necessary to combine several flaps.




Figure 3: 74 Year Old Woman with Lower Lip Carcinoma



Figure 4: Mc Gregor Unilateral Flap Postoperative Appearance after Four Years

The nasolabial flap is widely used to cover facial defects, including the reconstruction of postexcisional defects on the entire lower lips (Figure 5, 6).



Figure 5: Extensive Lower Lip Carcinoma in a 76 Year Old Man



Figure 6: Bilateral Nasolabial Flap Combined with Submental Advanced Flap

These flaps contained innervated muscle, has a good blood supply and motor function. The main disadvantage is the loss of saliva. The cross lip operation (Abbé flap) has an important role in improving the appearance of the lips by reducing tension and restoring lip balance, having the advantage of early sectioning of the pedicle and great flexibility, this flap can be used in combination with other flaps (Figure 7,8,9).



Figure 7: Extensive Lower Lip Carcinoma in a 52 year Old Man



Figure 8: Gillies Fan Flap Technique Combined With the Abbé Flap




Figure 9: Postoperative Appearance after Five Years

For tumors located in the central lower lip the Karapandzic flap, consisting in perioral advancement, remains a good option (Figure 10, 11).



Figure 10: Central Lower Lip Carcinoma in a 55 Year Old Man



Figure 11: Karapandzic Flap for Central Lip Reconstruction

The reconstruction of the commissure could be performed using a Brusari flap (Figure 12, 13, 14). Numerous techniques have been developed for lip reconstruction, the choice depends on the position of tumor and its extension level. Early stage tumors treated by direct suture have a good prognosis, compared with advanced lesions who need for reconstruction complex techniques under general anesthesia altering the appearance and the oral functionality.



Figure 12: Invasive Carcinoma at the Level of the Commissure in a 67 Year Old Patient



Figure 13: Intraoperative Aspect



Figure 14: Brusari Flap for Commissure Reconstruction.




Figure 15: Postoperative Appearance after Three Weeks.

At the time of diagnoses, 6 cases of patients with squamous cell carcinoma presented locoregional metastases. Recurrences after one year were recorded in 0.6 %. After 3 years the percentage increases to 1.8%. The survival rate at 3 years was 99%. Lymph node metastases reduce survival rate at 48%. Lymph node metastases appear to be more frequent in undifferentiated carcinoma than well differentiated tumors [10]. Recurring tumors and lymph node metastases is associated with a poor prognosis [11]. The prognosis is worse for tumours located at the commissure than the tumors from the middle part of the lip [12].

Conclusions

The interventions using local flaps with similar colour and texture provide the possibility of restoring aesthetic appearance with good functional results. The reconstruction after tumor excision with local flaps is a real challenge from the perspective of multiple surgical techniques using to restore all the structures in one session. The defect reconstruction for most extensive tumors, using local flaps, provides the possibility of restoring motor and sensory oral function. Postoperative wounds heal in maximum 3 weeks, and complication like necrosis, infection and hematoma are very rare. The patients with advanced tumors after wide excision and local plasty presented complication related to postoperative incontinence and incompetence, including vocalization, drooling and chewing disorder. This patients need kinetotherapy to improve lips mobility and flexibility and just a few patients return for commissuroplasty [13].

The incomplete excision, young patients, presence of other comorbidities and local recurrences are associated with poor prognosis. Increasing the resection margins according to the tumor size, decreases the risk of recurrence. Radiotherapy is indicated after the tumor resection. Lymph node neck metastases significantly reduces long term survival [14]. The diagnosis of these tumors is facilitated by a direct view of lesions and the local invasion should be avoided. For a plastic surgeon the aesthetic appearance has the same importance like the functional results.

Conflicts of interest statement: The authors declare no conflicts of interests.

References

1. Leiter U, Eigentler T, Garbe C (2014) Epidemiology of skin cancer. *Adv Exp Med Biol* 810: 120-40.
2. Vartanian JG, Carvalho AL, Araujo Filho, Hattori Junior M, Magrin J, et al. (2004) Predictive factors and distribution of lymph node metastasis in lip cancer patients and their implications on the treatment of the neck. *Oral Oncology* 40: 223-227.
3. Lim JL, Satem RS (2005) High levels of ultraviolet B exposure increase the risk of non-melanoma skin cancer in psoralen and ultraviolet A-treated patients. *J Invest Dermatol* 124: 505-13.
4. Moore SR, Johnson NW, Pierce AM, D Wilson (1999) The epidemiology of lip cancer: a review of global incidence and etiology. *Oral Dis* 5: 185-195.
5. Fabbrocini G, Triass M, Mauriello MC, Annunziata MC, De Vita V, et al. (2010) Epidemiology of skin cancer: role of some environmental factors. *Cancers* 2: 1980-9.
6. Papadopoulos O, Konofaos P, Tsantoulas Z, C Chrisostomidis, M Frangoulis et al. (2007) Lip defects due to tumor excision: Apropos of 899 cases. *Oral Oncology* 43: 204-212.
7. Baker SR. (1990) Current management of cancer of the lip. *Oncology* 4: 107-120.
8. Karapandzic M (1974) Reconstruction of lip defects by local arterial flaps. *Br J Plast Surg* 27: 93-97.
9. Gooris PJJ, Vermey A, Visscher JGAM, Fred R Burlage, Jan L N Roodenburg, et al. (2002) Supraomohyoid neck dissection in the management of cervical lymph node metastases of squamous cell carcinoma of the lower lip. *Head Neck* 24: 678-83.
10. Vukadinovic M, Jezdic Z, Petrovic M, Ljiljana M Medenica, Marko Lens, et al. (2007) Surgical management of squamous cell carcinoma of the lip: analysis of a 10-year experience in 223 patients. *J Oral Maxillofac Surg* 65: 675-679.
11. Bucur A, Stefanescu L (2004) Management of patients with squamous cell carcinoma of the lower lip and N0-neck. *J Craniomaxillofac Surg* 32: 16-21.
12. Visscher JGAM, Elsaker K, Grond AJK, J E van der Wal, I van der Waal, et al. (1998) Surgical treatment of squamous cell carcinoma of the lower lip: evaluation of long-term results and prognostic factors – A retrospective analysis of 184 patients. *J Oral Maxillofac Surg* 56: 814-820.
13. Zitsch RP (1993) Carcinoma of the lip. *Otolaryngol Clin North Am* 26: 265-277.
14. Zitsch RP, Lee BW, Smith RB (1999) Cervical lymph node metastases and squamous cell carcinoma of the lip. *Head Neck* 21: 447-453.